

## 炭酸カルシウム成長における不純物の影響についての理論的検討 Impurity effect for the formation mechanism of CaCO<sub>3</sub> polymorphs: ab initio study

川野 潤<sup>1\*</sup>, 佐久間 博<sup>2</sup>, 永井 隆哉<sup>3</sup>

Jun Kawano<sup>1\*</sup>, Hiroshi Sakuma<sup>2</sup>, Takaya Nagai<sup>3</sup>

<sup>1</sup>北海道大学・創成, <sup>2</sup>東京工業大学・院理工, <sup>3</sup>北海道大学・院理

<sup>1</sup>Hokkaido University, <sup>2</sup>Tokyo Institute of Technology, <sup>3</sup>Hokkaido University

The formation process of calcium carbonate polymorphs has been extensively investigated because of its importance in both geological and biological environments. Recently, it has been pointed out that the surface energy difference among polymorphs could determine their stability field, so that it becomes more important to analyze in detail their surface structures and the incorporation process of atoms and molecules into the growth surfaces.

In the present study, first-principles calculation of aragonite (001) surface was performed to theoretically analyze its surface structure and growth process. Especially, here we focused on the case in which Mg<sup>2+</sup> ion substitutes for Ca<sup>2+</sup> site as an impurity. Because, different from Ca<sup>2+</sup>, smaller Mg<sup>2+</sup> is unstable in the nine-fold cation positions of aragonite, it can hardly incorporate into bulk aragonite crystal. However it is expected to be able to substitute for Ca<sup>2+</sup> site in the surface bulk. Our simulation results show that Mg<sup>2+</sup> can incorporate into Ca<sup>2+</sup> sites in the surface with lower substitution energy than that for bulk, but Mg<sup>2+</sup> on the aragonite surface considerably affects the surface structure. This suggests the possibility that Mg<sup>2+</sup> could play an important role for aragonite formation and growth.

キーワード: aragonite, 表面構造, 第一原理計算

Keywords: aragonite, surface structure, first-principles calculation