

北西太平洋のマンガンクラストに記録されたミランコビッチ周期と環境変動 Milankovich cycle and environmental changes recorded in a ferromanganese crust from northwestern Pacific

小田 啓邦^{1*}, 宮城 磯治¹, 白井 朗²
Hirokuni Oda^{1*}, Isoji MIYAGI¹, Akira Usui²

¹ 産業技術総合研究所, ² 高知大学

¹National Institute of Advanced Industrial Science and Technology, ²Kochi University

A ferromanganese crust D96-m4, which was developed on the surface of basement rock, was taken from Shotoku seamount in the northwest Pacific. The magnetic fields of the thin sections were measured with the SQUID microscope. By correlating the polarity boundaries of the magnetic images (zero crossing lines) with the standard geomagnetic reversal timescale, the age of each zero crossing line was estimated and the growth rate was calculated as 5.1mm per million years (Oda et al., 2011). The beryllium isotope analysis of the same ferromanganese crust provided the growth rate estimate of 6.0 mm per million years, which is almost consistent with the estimate based on the magnetic method. Close investigation on the growth layers revealed the presence of Milankovich cycle for the last two million years including eccentricity (~100kyr) and obliquity (~40kyr). The identification of Milankovich cycle may provide high resolution age models of ferromanganese crusts. In the presentation, we will show the results of geochemical record as environmental changes measured by electron probe micro-analyzer on the crust based on the high resolution age model.

キーワード: 鉄マンガンクラスト, 北西太平洋, 古地磁気層序, ミランコビッチ周期, 成長縞, 化学組成変動

Keywords: ferromanganese crust, northwestern Pacific, magnetostratigraphy, Milankovich cycle, growth layer, geochemical variability