## Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



MZZ42-05

会場:104

時間:5月24日12:00-12:15

## 海洋大循環における北太平洋の役割について The Role of the North Pacific in the world ocean circulation

岡崎 裕典 <sup>1\*</sup> Yusuke Okazaki<sup>1\*</sup>

- 1 九州大学
- <sup>1</sup>Kyushu University

The North Pacific is known as a terminal region of Ocean Conveyer and abyssal circulation from the south flows into the North Pacific, and upwells to mid-depth and returns south as the Pacific Deep Water (Schmitz, 1996). No deep water forms in the North Pacific in the present ocean because the surface water of the North Pacific is not dense enough to sink into the deep basin (Warren, 1983). Instead, the North Pacific Intermediate Water (NPIW) originated from the Okhotsk Sea lies at depths of 300 to 800 m (Talley, 1993).

The glacial Pacific Ocean had two water masses: well-ventilated and nutrient-depleted glacial North Pacific Intermediate Water (GNPIW) above ~2000 m and less-ventilated and nutrient-enriched deep water below ~2000 m (Keigwin, 1998; Matsumoto et al., 2002). GNPIW is a thicker and more deeply penetrating water mass than the present NPIW. The possible source of GNPIW was possibly in the Bering Sea inferred from microfossil (Ohkushi et al., 2003) and neodymium isotope evidence (Horikawa et al., 2010). A switch of meridional overturning circulation between the North Atlantic and the North Pacific during the last glacial termination was suggested by a collaborative study of sedimentary proxy data and climate modeling (Okazaki et al., 2010).

The North Pacific appears to have played an active role in global ocean circulation, not always passive during the last glacial cycles. The Role of the North Pacific in the world ocean circulation of the past will be discussed.

## References:

Horikawa K, Asahara Y, Yamamoto K and Okazaki Y (2010) Intermediate water formation in the Bering Sea during glacial periods: Evidence from neodymium isotope ratios, Geology 38: 435-438.

Keigwin LD (1998) Glacial-age hydrography of the far northwest Pacific Ocean, Paleoceanography 13: 323-339.

Matsumoto K, Oba T, Lynch-Stieglitz J and Yamamoto H (2002) Interior hydrography and circulation of the glacial Pacific Ocean, Quaternary Science Reviews 21: 1693-1704.

Ohkushi K, Itaki T. and Nemoto N (2003) Last Glacial? Holocene change in intermediate-water ventilation in the Northwestern Pacific, Quaternary Science Reviews 22: 1477-1484.

Okazaki Y et al. (2010) Deepwater Formation in the North Pacific During the Last Glacial Termination, Science 329: 200-204. Schmitz WJ (1996) Woods Hole Oceanographic Institution Technical Report, WHOI-96-0B, 237 pp.

Talley LD (1993) Distribution and Formation of North Pacific Intermediate Water, Journal of Physical Oceanography 23: 517-537

Warren BA (1983) Why is no deep water formed in the North Pacific? Journal of Marine Research 41: 327-347.

キーワード: 北太平洋, 海洋大循環, 海洋ゲートウェイ, 北半球氷床発達