Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



会場:104



時間:5月24日12:30-12:45

## 人為起源ダスト増加による我が国での重金属・栄養塩負荷の増大に関する研究 Recent enrichment of nutrient and heavy metal deposition in Japanese mountain lakes due to anthropogenic dust

槻木 玲美<sup>1</sup>\*, 兵藤不二夫<sup>2</sup>, 加三千宣<sup>1</sup>, 牧野渡<sup>3</sup>, 谷幸則<sup>4</sup>, 占部城太郎<sup>3</sup> Narumi Tsugeki<sup>1</sup>\*, Fujio Hyodo<sup>2</sup>, Michinobu Kuwae<sup>1</sup>, Wataru Makino<sup>3</sup>, Yukinori Tani<sup>4</sup>, Jotaro Urabe<sup>3</sup>

## <sup>1</sup> 愛媛大学 上級研究員センタ ,<sup>2</sup> 岡山大学 異分野融合先端研究コア,<sup>3</sup> 東北大学大学院 生命科学研究科,<sup>4</sup> 静岡県立 大 環境科学研究所

<sup>1</sup>Senior Research Fellow Center, Ehime University, <sup>2</sup>Research Core for Interdisciplinary Sciences, Okayama University, <sup>3</sup>Graduate School of Life Science, Tohoku University, <sup>4</sup>Institute for Environmental Sciences, University of Shizuoka

Recent studies revealed that nutrient inputs such as nitrogen and phosphorus through atmospheric deposition to aquatic ecosystems have been increasing (Elser et al. 2009; Neff et al. 2008). In addition to nutrients, large emissions of heavy metals such as antimony (Sb) and indium (In) to the atmosphere is expected to be extensive (Filella et al., 2002; Tian et al., 2012; White and Hemond, 2012). East Asia plays an important role in global anthropogenic emissions, but little is known about the effects of nutrient emissions on terrestrial ecosystems and changes in the extent of the metal pollution during its rapid economic growth in recent decades. In this study, we examined fossil pigments and zooplankton remains in dated sediments taken from high mountain lakes at some Japanese National Parks in Hokkaido and Honshu area, to uncover the historical changes of plankton community over the past 100 years. Simultaneously, we measured the geochemical variables such as heavy metals, nitrogen and lead stable isotope to uncover the historical changes of metal deposition, and to identify causal factors including dust source regions. Sedimentary results showed that the fluxes of heavy elements of Sb and In increased at Lake Hachiman-Numa and Hourai-Numa in recent years (Kuwae et al. 2013). Furthermore, the fluxes of phytoplankton abundance in Lake Hourai-Numa (Tsugeki et al. 2012) and Lake Mikurigaike and Niseko-Onuma drastically increased since around 1990 when N stable isotope ratios in sediments decreased, probably due to expanding atmospheric N deposition. In parallel with this, Daphnia, a keystone herbivore, increased. During this period, there seems not to be expanding human activities in the watershed around these lakes, suggesting that the increases in nutrients and heavy metals were not resulted from inputs from watershed. Alternatively, Pb stable isotope data in Lake Hachiman-Numa and Hourai-Numa showed that dust deposition with nutrient and metal substances originated from the Asian continent were increasingly transported to study lakes in recent years (Tsugeki et al. 2012; Kuwae et al. 2013). These results imply that long-range transports of anthropogenic dusts have promoted not only Sb and In deposition but also eutrophication in a wide range of Japanese lakes even far from direct human disturbance.

Elser JJ, Andersen T, Baron JS, Bergstrom AK, Jansson M, Kyle M, Nydick KR, Steger L, Hessen DO (2009) Shifts in lake N: P stoichiometry and nutrient limitation driven by atmospheric nitrogen deposition. Science 326:835-837.

Filella M, Belzile N, Chen Y-W (2002) Antimony in the environment: a review focused on natural waters: I. Occurrence. Earth-Science Reviews 57:125-76.

Kuwae M, Tsugeki NK, Agusa T, Toyoda K, Tani Y, Ueda S, Tanabe S, and Urabe J (2013) Sedimentary records of metal depositions in Japanese alpine lakes for the last 250 years: Recent enrichments of airborne Sb and In in East Asia. Science of the Total Environment, 442: 189-197.

Neff JC, Ballantyne AP, Farmer GL, Mahowald NM, Conroy JL, Landry CC, Overpeck JT, Painter TH, Lawrence CR, Reynolds RL (2008) Increasing eolian dust deposition in the western United States linked to human activity. Nature Geo 1:189-195.

Tian H, Cheng K, Wang Y, Zhao D, Lu L, JiaW et al. (2012) Temporal and spatial variation characteristics of atmospheric emissions of Cd, Cr, and Pb from coal in China. Atmospheric Environment 50:157-63.

Tsugeki NK, Agusa T, Ueda S, Kuwae M, Oda H, Tanabe S, Tani Y, Toyoda K, Wang W, Urabe J (2012) Eutrophication of mountain lakes in Japan due to increasing deposition of anthropogenically-produced dust. Ecological Research 27: 1041-1052.

White SJO, Hemond HF (2012) The anthrobiogeochemical cycle of indium: a review of the natural and anthropogenic cycling of indium in the environment. Critical Reviews in Environmental Science and Technology 42:155-186.

キーワード:古陸水学,アジア大陸,人為起源ダストと栄養塩負荷,プランクトン,重金属,安定同位体

Keywords: paleolimnology, Asian continent, anthropogenic dust and nutrient deposition, plankton, heavy metal, stable isotope