

## 高緯度サンゴを用いた、酸素同位体比と Sr/Ca 比の骨格成長量依存性の評価 Estimation of the growth-rate influences on the oxygen isotope and Sr/Ca ratios in the Porites at high latitudes

平林 頌子<sup>1\*</sup>, 横山 祐典<sup>1</sup>, 鈴木 淳<sup>2</sup>, 川久保 友太<sup>1</sup>, 宮入 陽介<sup>1</sup>, 岡井 貴司<sup>2</sup>, 野島 哲<sup>3</sup>

Shoko Hirabayashi<sup>1\*</sup>, Yusuke Yokoyama<sup>1</sup>, Atsushi Suzuki<sup>2</sup>, KAWAKUBO, Yuta<sup>1</sup>, Yosuke Miyairi<sup>1</sup>, Takashi Okai<sup>2</sup>, NOJIMA, Satoshi<sup>3</sup>

<sup>1</sup> 東京大学大気海洋研究所, <sup>2</sup> 独立行政法人 産業技術総合研究所地質情報研究部門, <sup>3</sup> 九州大学理学部附属天草臨海実験所  
<sup>1</sup> Atmosphere and Ocean Research Institute, The University of Tokyo, <sup>2</sup> Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology (AIST), <sup>3</sup> Amakusa Marine Biological Laboratory, Kyushu University

The skeletal oxygen isotope in the coral is used as a good proxy to reconstruct sea surface temperature (SST). On the other hand, some studies showed that coral growth rates may affect the oxygen isotope in coral (McConnaughey 1989a; Felis et al., 2003; Suzuki et al., 2005; Hayashi et al., 2013). Here we reanalyzed the same Porites coral in Omata et al. (2006), which was collected in the eastern of Ushibuka in the Amakusa area, Japan, which is located at slightly lower latitude than the northern limit of the hermatypic corals. We measured Sr/Ca ratio and reconstructed SST. Compared to the oxygen isotope ratio measured by Omata et al. (2006), our results showed that Sr/Ca ratio is the robust SST proxy which is independent of its growth rate. We suggest that Sr/Ca ratio is more suitable for reconstruction of SST using the small growth-rate Porites corals, especially in high latitudes.

キーワード: サンゴ骨格中の酸素同位体, Sr/Ca 比, 高緯度, サンゴの成長率

Keywords: skeletal oxygen isotope in the coral, Sr/Ca ratio, high latitudes, coral growth rates