Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.

会場:101B

PEM08-11



時間:5月20日17:00-17:15

Preliminary result of the comparison between hi-resolution TEC map and airglow images observed by ISS/IMAP-VISI Preliminary result of the comparison between high-resolution TEC map and airglow im-

## ages observed by ISS/IMAP-VISI

西岡 未知<sup>1\*</sup>, 津川 卓也<sup>1</sup>, 齊藤 昭則<sup>2</sup>, 坂野井 健<sup>4</sup>, 秋谷 祐亮<sup>2</sup>, 大塚 雄一<sup>3</sup>, 石井 守<sup>1</sup> Michi Nishioka<sup>1\*</sup>, Takuya Tsugawa<sup>1</sup>, Akinori Saito<sup>2</sup>, Takeshi Sakanoi<sup>4</sup>, Yusuke Akiya<sup>2</sup>, Yuichi Otsuka<sup>3</sup>, Mamoru Ishii<sup>1</sup>

<sup>1</sup>(独)情報通信研究機構,<sup>2</sup>京都大学大学院理学研究科,<sup>3</sup>東北大学大学院理学研究科,<sup>4</sup>名古屋大学太陽地球環境研究所 <sup>1</sup>National Institute of Information and Communications Technology, <sup>2</sup>Graduate school of science, Kyoto University, <sup>3</sup>Graduate school of science, Tohoku University, <sup>4</sup>Solar-Terrestrial Environment Laboratory, Nagoya University

Two-dimensional Total Electron Content (TEC) map has been used for revealing characteristics of meso-scale ionospheric disturbances, such as Equatorial Plasma Bubble (EPB) and Travelling Ionospheric Disturbances (TID). We have developed high-resolution TEC map and started "Dense Regional And Worldwide International Networks of GNSS-TEC observation (DRAWING-TEC) " project (http://seg-web.nict.go.jp/GPS /DRAWING-TEC/). The DRAWING-TEC project aims to expand the high-resolution TEC observation area by sharing TEC data in the newly standardized GTEX format. The project also provides regional and global maps of absolute value of TEC, de-trended TEC with 60-min window, and Rate of TEC change Index (ROTI). As of 2013, regional maps are produced over North America and Europe in addition to Japan. The spatial resolution of TEC maps of North America and Europe is 0.15 deg x 0.15 deg with a 5x5 pixel smoothing. The temporal resolution of these maps is 30 seconds. These regional maps are powerful tools for studying characteristics of EPBs and TIDs. We compared the regional TEC map with the 630 nm airglow image obtained by ISS-IMAP/VISI. The preliminary results of the comparison will be shown in the presentation.

キーワード: 電離圏全電子数, 宇宙ステーション, 伝搬性電離圏擾乱, 赤道域プラズマバブル Keywords: Total Electron Content map, ISS/IMAP, traveling ionospheric disturbance, equatorial plasma bubble