Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



PEM09-12

会場:101B

関東地方における VLF 帯雷電磁波観測データを用いた落雷規模の推定 Estimation of lightning magnitude using VLF sferics in Kanto rigion

工藤 剛史^{1*}, 高橋 幸弘¹, 佐藤 光輝¹, 長 康平²

Takeshi Kudo1*, Yukihiro Takahashi1, Mitsuteru Sato1, Kohei Osa2

1北海道大学大学院理学院宇宙理学専攻,2株式会社ウェザーニューズ

¹Department of Cosmosciences, Graduate school of Science, Hokkaido University, ²WEATHERNEWS INC.

It is pointed out that the relationship between atmospheric parameters and occurrence frequency of lightning is closely correlated. By the current lightning detection systems operated in Japan, which measure higher frequency than VLF range, only the information of peak current of the stroke is estimated and information of its magnitude, such as charge moment change (CMC) cannot be derived. Since the most of the electromagnetic energy of lightning concentrates in the frequency range less than 10 kHz, in order to estimate CMC of lightning stroke, we need to observe the electromagnetic waves in lower frequency range than existing lightning networks which make use of order of several 10s of kHz to M Hz range. Carrying out the continuous observation of VLF sferic, we will investigate the quantitative relationship between atmospheric parameters and lightning activity including information of electrical magnitude for the first time.

We are constructing a VLF observation network with identical observation system, consisting of three observation sites in Kanto region, which enables us to geolocate lightning stroke by time of arrival (TOA) or direction finding methods with an error of 10 km. Each observation system is composed of two horizontal magnetic loop antennas and a vertical electric dipole antenna, receiver, PC and GPS clock. The higher cutoff and sampling frequency are 40 kHz and 80 kHz, respectively.

In order to estimate CMC, the temporal variation of source current with peak intensity is required. We estimated the peak current using a method proposed by Yanagi [2012]. The temporal variation of source current was estimated from the groundwave of electric field, using amplitude and zero-crossing time of the waveform.

In this presentation, we will report the situation of construction of this observation network and the estimation methodology of lightning peak current using sferic data. Also preliminary results based on the data obtained at Yamanashi station are introduced. This research is supported by grant in aid by Kisho-Bunka-Souzou center and Kakenhi (Kiban-A) No. 2425300202.

Yanagi, Y., Development of VLF electromagnetic wave observation system and estimation method of lightning peak current, Master thesis, Hokkaido University, 2012.

Keywords: lightning, charge moment change