Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



## 会場:202

時間:5月23日10:30-10:45

## ー点観測を用いた太陽風磁気フラックスロープの時間発展の再現 Reconstruction of an evolving magnetic flux rope in the solar wind from single-spacecraft data

長谷川 洋<sup>1\*</sup>, Bengt Sonnerup<sup>2</sup>, Qiang Hu<sup>3</sup>, 中村 琢磨<sup>4</sup> Hiroshi Hasegawa<sup>1\*</sup>, Bengt Sonnerup<sup>2</sup>, Qiang Hu<sup>3</sup>, Takuma Nakamura<sup>4</sup>

<sup>1</sup>JAXA 宇宙科学研究所,<sup>2</sup>ダートマス大学,<sup>3</sup>アラバマ大学,<sup>4</sup>ロスアラモス国立研究所

<sup>1</sup>Institute of Space and Astronautical Science, JAXA, <sup>2</sup>Dartmouth College, <sup>3</sup>University of Alabama in Huntsville, <sup>4</sup>Los Alamos National Laboratory

In situ measurements, often made by single spacecraft, have a difficulty in revealing spatiotemporal evolution of space plasma structures. We present a single-spacecraft method for decomposing spatial and temporal variations of physical quantities at points along the path of a spacecraft in spacetime, which can be used for reconstruction of slow evolution of two-dimensional (2D) and magneto-hydrostatic structures (namely, Grad-Shafranov equilibria) (Sonnerup and Hasegawa, 2010). The method is applicable to structures that are in Grad-Shafranov equilibrium and in which the flow is incompressible and the frozen-in condition is satisfied. Benchmark tests are conducted by use of synthetic data taken by a virtual spacecraft that traverses, at a constant velocity, a magnetic flux rope growing in a 2D magnetohydrodynamic simulation of magnetic reconnection. It is demonstrated that the new method can better recover the quantities in spacetime than does an earlier version, in which time aliasing effects had not been removed (Hasegawa et al., 2010). The application to a flux rope observed on 25-26 March 1998 by the ACE spacecraft in the solar wind suggests that its core part was evolving in an intriguing way during the ~17 hour interval of traversal.

Sonnerup, B. U. O., and H. Hasegawa (2010), On slowly evolving Grad-Shafranov equilibria, J. Geophys. Res., 115, A11218, doi:10.1029/2010JA015678.

Hasegawa, H., B. U. O. Sonnerup, and T. Nakamura (2010), Recovery of time evolution of Grad-Shafranov equilibria from single-spacecraft data: Benchmarking and application to a flux transfer event, J. Geophys. Res., 115, A11219, doi:10.1029/2010JA015679.

キーワード: 太陽風, 磁気フラックスロープ, グラッド・シャフラノフ方程式, 電磁静水圧平衡 Keywords: solar wind, magnetic flux rope, Grad-Shafranov equation, magnetohydrostatic equilibrium