

サブストーム発生時に於けるオーロラビーズの発生と地磁気脈動Piとの関係 Relationships between auroral beads evolution and Pi pulsations at substorm onset

櫻井 亨^{1*}, 元場 哲郎², 細川 敬祐³, 門倉 昭⁴, 田中 良昌⁵, 宮岡 宏⁶, 佐藤 夏雄⁷

Tohru Sakurai^{1*}, Tetsuo Motoba², Keisuke Hosokawa³, Akira Kadokura⁴, Yoshimasa Tanaka⁵, Hiroshi Miyaoka⁶, Natsuo Sato⁷

¹ 東海大学, ² ジョンホプキンス大学, ³ 電気通信大学, ⁴ 国立極地研究所, ⁵ 国立研究所, ⁶ 国立極地研究所, ⁷ 国立極地研究所
¹Tokai University, ²John Hopkins University Applied Physics Laboratory, ³University of Electro-Communications, ⁴National Institute Polar Research, ⁵National Institute Polar Research, ⁶National Institute Polar Research, ⁷National Institute Polar Research

Auroral beads often appeared during presubstorm onset phase, which has been considered as one of important auroral phenomena relating to substorm expansion onset. Recently Motoba et al.,(2012) have reported optical signatures of auroral beads evolution in all ? sky images observed at a conjugate pair ? station, Syowa in Antarctica and Tjornes in Iceland, in which auroral beads appeared almost at the same time, as an inter ? hemispheric signature and the evolution was stable in the first stage, and then developed to a larger scale spiral form (undulations), at the same time at both the conjugate stations. These interhemispheric similarities suggest that there must be a common driver in the magnetotail equatorial region. On the other hand, it has been well known that Pi pulsation is another important signature at substorm onset. In this paper we examined relationships between auroral beads evolution and Pi pulsation onsets during a few minutes before the substorm expansion onset for the 30 September 2012 substorm event. Results show that there exist very good relationships between Pi pulsation onsets and auroral beads evolutions, particularly on the signatures for auroral beads sudden evolution to spiral structures and/or undulations, which was coincident to appearance of Pi 2 pulsation, and auroral beads brightness enhancement, which accompanied Pi 1 pulsation onset. The appearances of these Pi 1 and Pi 2 pulsations show a good conjugacy. These results show very good one-to-one correspondence between auroral beads dynamics and Pi 1 and 2 onsets, suggesting a common physical mechanism controlling their dynamics in the magnetotail equatorial region.

Keywords: substorm, aurora, pulsations