Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.

PPS01-04

会場:201B



時間:5月21日15:00-15:15

## 非熱的中性粒子観測器による木星探査 Measurement low-energy energetic neutral atoms around Jupiter's satellite

浅村 和史 <sup>1</sup>\*, 二穴喜文 <sup>2</sup>, 三好 由純 <sup>3</sup>, 坂野井 健 <sup>4</sup>, PEP チーム <sup>2</sup> Kazushi Asamura<sup>1</sup>\*, Yoshifumi Futaana<sup>2</sup>, Yoshizumi Miyoshi<sup>3</sup>, Takeshi Sakanoi<sup>4</sup>, PEP team<sup>2</sup>

<sup>1</sup>ISAS/JAXA, <sup>2</sup>IRF, Sweden, <sup>3</sup>名古屋大, <sup>4</sup>東北大 <sup>1</sup>ISAS/JAXA, <sup>2</sup>IRF, Sweden, <sup>3</sup>Nagoya U., <sup>4</sup>Tohoku U.

We are proposing a measurement of low-energy (10eV-3keV) energetic neutral atoms around Jupiter's satellite especially Ganymede. Ganymede has its own intrinsic magnetic moment. There is considered to be a mini-magnetosphere around Ganymede because of interactions between plasma in Jovian magnetosphere and Ganmede's magnetic field. However, its characteristics will be different from terrstrical one, since Alfven mach number of upstream plasma flow (corotational plasma flow around Jupiter) is small. JNA (Jovian Neutral Analyzer) will reveal characteristics of Ganymede's magnetosphere in terms of measurement of scattered/sputtered particles generated by precipitation of plasma particles onto Ganymede's surface. Measurement of these particles will provide spatial distribution of plasmas in remote sense, since electric/magnetic field do not affect trajectories of neutral particles. JNA is a part of PEP (Particle Environment Package) led by Swedish Institute of Space Physics, which was proposed as potential instruments onboard JUICE mission.

We will discuss current status of JNA.

Keywords: low-energy energetic neutral atoms, Ganymede, magnetosphere, remote observation