

## Evidence for a dynamic event recorded in HED meteorites

Masaaki Miyahara<sup>1\*</sup>, Eiji Ohtani<sup>1</sup>, Akira Yamaguchi<sup>2</sup>, Shin Ozawa<sup>2</sup>, Takeshi Sakai<sup>3</sup>, Hiraon Naohisa<sup>4</sup>

<sup>1</sup>Graduate School of Science, Tohoku University, <sup>2</sup>National Institute of Polar Research, <sup>3</sup>Geodynamics Research Center, Ehime University, <sup>4</sup>Japan Synchrotron Radiation Research Institute

It is expected that HED meteorites originate from one of the largest asteroids in the solar system, 4 Vesta. Recent Dawn mission operated by NASA also supports the prediction [1][2]. Dawn mission clearly revealed the existence of many craters on 4 Vesta, which are the records of heavy meteorite bombardments. The existence of a high-pressure polymorph in a shocked meteorite is a robust evidence for a dynamic event on its parent-body (e.g., [3][4]). A high-pressure polymorph can be used for estimating the magnitude of a dynamic event (e.g., [5][6]). Some previous studies propose that 4 Vesta might suffer from late heavy bombardment (LHB) as well as the moon [7]. However, a high-pressure polymorph has not been found in HED meteorite so far.

We got one of eucrite samples, Bereba to clarify a dynamic event occurred on 4 Vesta using a high-pressure polymorph. Present Bereba sample has several shock-melt veins, implying that it was heavily shocked. Accordingly, we investigated Bereba using Raman spectroscopy, FEG-SEM, synchrotron(s)-XRD and FIB-TEM techniques to clarify a record of a dynamic event occurred on 4 Vesta.

We focused our interest on silica phase in this study. Raman spectroscopy analyses show that the silica grains in the host-rock of Bereba are cristobalite, tridymite and quartz. Most quartz grains entrained in the shock-melt veins are partly replaced with coesite. BSE images show that silica grains entrained in or adjacent to the shock-melt veins have network-like and/or lamellae-like textures. Raman spectroscopy, s-XRD analyses and TEM images indicate that such silica grains include coesite, stishovite and silica glass along with quartz.

We found the high-pressure polymorphs of silica from HED meteorites for the first time. The existence of stishovite indicates that pressure condition recoded in Bereba should be ~8 GPa at least based on a phase diagram obtained from static high-pressure and high-temperature synthetic experiments [8]. Two giant impact basins on 4 Vesta are depicted by Dawn mission. Crater chronology obtained by Dawn mission reveals that the giant impact basins were formed around 1.0 Ga ago [9]. Its fragments became Vesta family in asteroid belt, and a part of them fell into the Earth as HED meteorites. The existence of high-pressure polymorphs in Bereba may support the giant impact occurred on 4 Vesta

### References:

- [1] De Sanctis M.C. et al. (2012) *Science*, 336, 697-700. [2] Russell C.T. et al. (2012) *Science*, 336, 684-686. [3] Ohtani E. et al. (2011) *PNAS*, 108, 463-466. [4] Miyahara M. et al. (2011) *PNAS*, 108, 5999-6003. [5] Chen M. et al. (1996) *Science*, 271, 1570-1573. [6] Ohtani E. et al. (2004) *EPSL*, 227, 505-515. [7] Bogard D.D. (2011) *Chem. Erde.*, 71, 207-226. [8] Zhang J.R.C. et al. (1993) *J. Geophys. Res.* 98, 19785-19793. [9] Marchi S. et al. (2012) *Science*, 336, 690-694.