

炭素循環の機能する系外地球型水惑星の表層環境と進化 Climate evolution of extrasolar terrestrial planets with water and carbonate-silicate geochemical cycle

門屋 辰太郎^{1*}, 田近 英一²
Shintaro Kadoya^{1*}, Eiichi Tajika²

¹ 東京大学大学院理学系研究科, ² 東京大学大学院新領域創成科学研究科
¹The University of Tokyo, ²The University of Tokyo

The surface environment of a habitable planet has been constrained in terms of orbital semi-major axis and an effective solar flux, focusing on the existence of liquid water, that is, the habitable zone (HZ) around main sequence stars (e.g., Kasting et al., 1993). It has been also pointed out that the carbonate-silicate geochemical cycle would be essential for maintaining the climate of a habitable planet (e.g., Tajika, 2003). However, the whole picture of evolution of climate of planets with carbonate-silicate geochemical cycle has not been known.

In this study, we investigate the climate evolution of an Earth-like planet (actually, the Earth itself) around a G-type star (= Sun). Steady states of climate of Earth-like planets are estimated systematically with a simple climate model coupled with a carbonate-silicate geochemical cycle model. We classified climates of Earth-like planets within the HZ into three modes, in terms of stabilizing mechanism. Then, the climate evolution is estimated based on the steady state solutions of the climate with models of the stellar evolution and the thermal evolution of planetary interiors. The results indicate that, on an Earth-like planet (the size of the Earth) orbiting around a G-Type star, climate depends on the thermal history of the planet in the early stages of its lifetime, and then depends on the stellar evolution. The climate evolutions are also estimated for the different mass both for stars and planets.

キーワード: 太陽系外惑星系, ハビタブルゾーン, ハビタブルプラネット, 炭素循環, EBM

Keywords: Extrasolar planetary system, Habitable zone, Habitable planet, Carbonate-silicate geochemical cycle, EBM