## Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



PPS23-06

会場:201A

時間:5月23日10:15-10:30

## 月面の二次クレーター検出のためのクレーター空間分布の評価方法の開発 Development of evaluating method of Spatial distribution of craters on Lunar Surface for detection of secondary craters

木下 達生  $^{1*}$ , 本田 親寿  $^{1}$ , 平田 成  $^{1}$ , 諸田 智克  $^{2}$  Tatsuo Kinoshita $^{1*}$ , Chikatoshi Honda $^{1}$ , Naru Hirata $^{1}$ , Tomokatsu Morota $^{2}$ 

1 会津大学先端情報科学研究センター宇宙情報科学クラスタ, 2 名古屋大学大学院環境学研究科

Secondary craters are impact craters formed by ejecta blocks that were thrown out of a primary crater formation. The secondary craters give a biased spatial distribution of craters. For a crater chronology, researchers extract the secondary craters from the surface including primary and secondary craters based on his or her subjective views.

The purpose of this research is to develop an algorithm for evaluating spatial distribution of craters on lunar and planetary images. In our algorithm, clustering analysis (S-LINK, Group average etc.) applies to ideal spatial distribution of craters and observed spatial distribution of craters, and evaluates whether a non-random portion in obtained image by comparing clustering analysis results of ideal and observed craters. We demonstrated for two regions in Mare Crisium and some Apollo landing sites. As a result, most of clustered secondary craters are detected quantitatively by our algorithm.

キーワード: 月, 二次クレーター, クラスター分析 Keywords: Moon, Secondary crater, Cluster analysis

<sup>&</sup>lt;sup>1</sup>CAIST Arc-Space, University of Aizu, <sup>2</sup>Graduate School of Environmental Studies, Nagoya University