

微隕石研究のための新しい分析スキームの開発と特異な鉱物学的特徴を持つ微隕石の発見

Development of a new analytical scheme for micrometeorites and discovery of micrometeorites with intriguing mineralogy

辻本真一¹, 野口 高明^{1*}, 岡崎 隆司², 中村 智樹³, 海老原 充⁴, 伊藤 正一⁵, 永原 裕子⁶, 橘 省吾⁵, 寺田 健太郎⁷, 藪田 ひかる⁷
Shinichi Tsujimoto¹, Takaaki Noguchi^{1*}, Ryuji Okazaki², Tomoki Nakamura³, Mitsuru Ebihara⁴, Shoichi Itoh⁵, Hiroko Nagahara⁶, Shogo Tachibana⁵, Kentaro Terada⁷, Hikaru Yabuta⁷

¹ 茨城大学, ² 九州大学, ³ 東北大学, ⁴ 首都大学東京, ⁵ 北海道大学, ⁶ 東京大学, ⁷ 大阪大学

¹Ibaraki University, ²Kyushu University, ³Tohoku University, ⁴Tokyo Metropolitan University, ⁵Hokkaido University, ⁶University of Tokyo, ⁷Osaka University

Introduction: Combined mineralogical and isotopic studies of individual Wild 2 cometary particles revealed that the materials formed in the inner solar system had been transported to the outer solar system before the formation of the Wild 2 comet [e. g. 1,2,3,4]. We sought another analytical scheme for MM (micrometeorite) studies. In this new analytical scheme, we have two objectives: identification of nonchondritic MMs and identification of asteroidal MMs with mineralogy indistinguishable from CP IDPs, which are regarded as cometary grains [5].

Samples and methods: MMs used in this study were found in the surface snow collected near the Dome Fuji Station, Antarctica in 2003 and 2010. The surface snow was melted and filtered in a clear room. After identification of MMs, we performed SR-XRD, FIB section preparation, TEM, micro-Raman, SEM, EPMA, INAA, and/or noble gas mass spectroscopy for each MM.

Results and discussion: We investigated twelve MMs and could classify them into five types based on their mineralogy: refractory MMs, chondrule-like MMs, fine-grained polycrystalline MMs, coarse-grained crystalline MMs, and phyllosilicate-rich MMs. Here we show the mineralogical results of two MMs with intriguing mineralogy.

Refractory MM A MM is composed mainly of anorthite, diopside, and spinel. A BSE image of the cross section of the MM showed that it has an amoeboid structure, in which small (<5 micrometer) Al-rich and Ti-bearing diopside exists on its surface and its interior and anorthite fills the interstices of diopside. The major minerals are similar to those in type C CAIs [6]. Although its amoeboid structure suggests low degrees of melting during the formation event, the MM has a compact interior, in which diopside and anorthite have triple junctions with $\sim 120^\circ$ angles. Therefore, the refractory object was probably formed by low degrees of melting and subsequent prolonged cooling. It is different from meteoritic type C CAIs that experienced intense melting and crystallization from melt droplets. Fine-grained material attached on the refractory object is composed of GEMS (glass with embedded metal and sulfide)-like objects, olivine, pyrrhotite, and carbonaceous material. Because GEMS-like objects have not been identified among meteorites yet, this MM may have derived from a comet.

Fine-grained polycrystalline MM A MM is composed of small (<400 nm) crystals of Fe-bearing olivine, Fe-free low-Ca pyroxene, Fe-Ni metal, Fe sulfide, amorphous silicate, and interstitial carbonaceous material. Although these crystals have often triple junctions with $\sim 120^\circ$ angles suggestive of recrystallization, both olivine and low-Ca pyroxene show almost no compositional zoning. The low-Ca pyroxene crystals elongate near the a-axis direction and are composed of a unit cell-order mixture of ortho and clino low-Ca pyroxene with stacking disorders. Its microstructure indicates rapid cooling ($>20\text{-}30\text{ K hr}^{-1}$) from above 1275K [7]. Both olivine and low-Ca pyroxene in this MM contain abundant tracks with $\sim 5 \times 10^{10}\text{ cm}^{-2}$, which corresponds to $\sim 10^4$ -year exposure to the solar wind [8]. Tracks in olivine are erased by flash heating above $\sim 600^\circ\text{C}$ [9]. Because typical cometary IDPs are heated to $>720^\circ\text{C}$ [10], it is plausible that the MM was derived from an asteroid instead of a comet although any materials similar to this MM have not been found among meteorites.

References: [1] Zolensky et al. (2006) Science 314, 1735-1739. [2] McKeegan et al. (2006) Science 314, 1724-1728. [3] Nakamura et al. (2008) Science 321, 1664-1667. [4] Matzel et al. (2010) Science 328, 483-486. [5] Ishii et al. (2008) Science 319, 447-450. [6] Krot et al. (2008) GCA72, 2534-2555. [7] Brearley and Jones (1993) LPS XXIV, 185-186. [8] Bradley (2006) In: Meteorites, Comets, and Planets. [9] Fraundorf et al. (1982) GCA 45, 915-943. [10] Joswiak et al. (2005) Workshop on dust in planetary systems (abstract), #4106.

キーワード: 微隕石, 透過電子顕微鏡, 希ガス, 集束イオンビーム, 放射化分析

Keywords: micrometeorites, TEM, noble gas, FIB, INAA