

ジュラ紀付加体に産する炭酸マンガンノジュール形成過程の地球化学的制約 A geochemical constraint on the formation process of a manganese carbonate nodule in the siliceous mudstone of the Juras

中田 亮一^{1*}, 白井孝明², 高橋 聡³, 鈴木 紀毅², 小川和広², 高橋 嘉夫¹

Ryoichi Nakada^{1*}, Takaaki Shirai², Satoshi Takahashi³, Noritoshi Suzuki², Kazuhiro Ogawa², Yoshio Takahashi¹

¹ 広島大・院理, ² 東北大・院理, ³ 東京大・理

¹Hiroshima Univ., ²Tohoku Univ., ³Univ. of Tokyo

Manganese (Mn) carbonate nodules, which are different from seafloor Mn nodules mainly composed of MnO₂, are occasionally embedded as a lens shape in the Jurassic accretionary complexes such as the Tamba-Mino-Ashio Belt in Japan. The interpretation of the formation process of Mn carbonate is still controversial, namely whether the Mn carbonate was formed primarily or secondarily. The optical observation of thin sections shows that Mn carbonate nodules are comprised with abraded grains of rhodochrosite spherule with radiolarians and are sedimentarily embedded in siliceous mudstone. Microfossil radiolarians from the Mn carbonate nodules and the host red siliceous mudstone are dated as the Bajocian (170.3 ? 168.3 million years ago), but radiolarians in the nodules are somewhat older than those in the host red siliceous mudstone. In this study, geochemical analyses focusing on the rare earth element (REE) were performed to unlock the formation process of Mn carbonate nodules in the Tamba-Mino-Ashio Belt.

Our analysis using the X-ray absorption near-edge structure on Ce shows dominance of trivalent Ce at present, despite of a positive Ce anomaly in the PAAS-normalized REE pattern of Mn carbonate. The REE adsorption experiment on synthesized MnCO₃ does not show any distinctive positive Ce anomaly and a thermodynamic calculation suggests the possible coexistence of authigenic rhodochrosite and spontaneous oxidation of Ce. The leaching experiment that can selectively decompose carbonate phase demonstrated no Ce anomaly in the carbonate phase of Mn carbonate and poor contribution to the bulk REE concentration. The most plausible account of all the observational and experimental results is that rhodochrosite grains were primarily formed on the depositional site and subsequently transferred to the different site where siliceous mudstone was deposited.

キーワード: 炭酸マンガン, 希土類元素, 放散虫, XAFS

Keywords: manganese carbonate, rare earth element, radiolarian, XAFS