

## 初期地球におけるコアへのカリウム・親石元素の分配と地球熱史・コア軽元素への制約

### Metal-silicate partitioning of lithophile elements and implications for potassium and light elements in the core

野村 龍一<sup>1\*</sup>, 廣瀬 敬<sup>2</sup>

Ryuichi Nomura<sup>1\*</sup>, Kei Hirose<sup>2</sup>

<sup>1</sup> 東京工業大学, <sup>2</sup> 東京工業大学 地球生命研究所, J A M S T E C

<sup>1</sup>Tokyo Institute of Technology, <sup>2</sup>ELSI, Tokyo Institute of Technology, JAMSTEC

The partitioning of lithophile elements (K, O, Si, Mg, Al, and Ca) between liquid metal and silicate melt were investigated up to 138 GPa and 5450 K in S-free/S-bearing Fe + K-doped pyrolite system, in order to constrain the amounts of radioactive <sup>40</sup>K and other light elements in the core. The obtained iron-potassium exchange coefficients show strong temperature (T) dependence but negligible effects of pressure (P) and sulfur content, not supporting the transition-metal-like behavior of potassium at high pressure. As a consequence, present experiments suggest only ~10 ppm potassium in the core, which yields present-day heat production of ~0.1 TW, even when we assume the entire core-mantle chemical equilibrium at 136 GPa and 5300 K (liquidus of pyrolitic mantle). On the other hand, the core dissolves substantial amounts of silicon and oxygen as a consequence of reaction with a basal magma ocean at 4500-5000 K, which account for the 10% core density deficit. In addition, quenched liquid iron obtained in relatively high-temperature experiments included certain amounts of Mg, Al, and Ca, suggesting that these elements may have been once incorporated into the core at the time of giant impact. A more realistic model for the Earth's core must consider the combined effects of material that equilibrated at modest P-T with the material that was added at very high P-T, and the effect of latter components are focused in this study.

Keywords: metal-silicate partitioning, high pressure, core formation, magma ocean, potassium, light elements