

K-Ar年代測定における試料サイズと過剰アルゴンの関係 Reduction of extraneous ^{40}Ar contamination for accurate K-Ar age determinations: an experimental study in various sample

山崎 誠子^{1*}, 磯部 翔太郎², 佐藤 洋樹², 田上 高広²
Seiko Yamasaki^{1*}, Shotaro Isobe², Hiroki Sato², Takahiro Tagami²

¹ 産業技術総合研究所, ² 京都大学
¹Geological Survey of Japan, AIST, ²Kyoto University

A fundamental assumption of K-Ar dating is that the samples initially contained no radiogenic ^{40}Ar , but sometimes rocks contain radiogenic ^{40}Ar called extraneous ^{40}Ar . Some previous study reported argon isotopes of historical lavas had anomalously high $^{40}\text{Ar}/^{36}\text{Ar}$ ratios, and show old apparent ages. Since extraneous ^{40}Ar is likely contained in the phenocrysts and xenoliths, groundmass samples are generally prepared for analysis. Besides, Ozawa et al. (2005) showed fine-grained groundmass samples had less extraneous ^{40}Ar contamination, and suggested that extraneous ^{40}Ar is contained in fluid inclusions or vesicles and released during crushing. We measure argon isotopic ratios in various sizes of young lava samples, and investigated the reduction of extraneous ^{40}Ar contamination. The finer samples roughly showed lower $^{40}\text{Ar}/^{36}\text{Ar}$ ratios but more difficult to handling of the preparation such as mineral separation and wrapping in foils for isotopic measurements.

キーワード: K-Ar年代測定, 過剰 ^{40}Ar , 粒度
Keywords: K-Ar dating, extraneous ^{40}Ar , sample size