

最上部マントルにおける MORB 生成の多様性: 北部オマーンオフィオライトを例として
Diversity of MORB genesis within the uppermost mantle: an example from the northern Oman ophiolite

秋澤 紀克¹, 荒井 章司^{1*}, 田村 明弘¹, 小澤 一仁²
Norikatsu Akizawa¹, Shoji Arai^{1*}, Akihiro Tamura¹, Kazuhito Ozawa²

¹ 金沢大学・地球, ² 東京大学・地球惑星環境学科
¹Earth Sciences, Kanazawa Univ., ²Earth and Planetary Science

Dunite bands and veins in the ophiolitic mantle peridotite are interpreted as fossil melt conduits within the suboceanic mantle. In particular, concordant dunite bands are possibly important as the melt conduits through which parental melts of MORB (mid-ocean ridge basalts) were transported to shallower mantle beneath the ridge axis. However, no detailed petrological data of concordant dunite bands and surrounding peridotites have been published. We conducted sampling of concordant dunite bands and its aureole from various "stratigraphic levels" in the mantle section from an estimated ancient-segment center and its end in the northern Oman ophiolite. They are various both in thickness (few millimeters to few meters) and in frequency of appearance. Dunite bands are almost pyroxene-free, and their orthopyroxenes, if any, are vermicular in shape.

Mineral chemistry shows systematic variations in the wall peridotites toward the dunite bands: (1) a decrease in Fo content (92 to 90.5) of olivines, (2) an increase in Cr/(Cr + Al) atomic ratio (0.5 to 0.6) and TiO₂ content (nil to 0.25 wt %) in spinels, and (3) an increase in Na₂O content (almost nil to 0.2 wt%) of clinopyroxene. In ambient residual peridotites, rare earth element (REE) patterns of clinopyroxene incline from light-REE (LREE) to heavy-REE (HREE) monotonously. The REE pattern of clinopyroxene in dunitites and surrounding peridotites show various shapes, depending on the position, the segment center to end: gentle slope from HREE to LREE at the segment center, and U-shaped at the segment end.

We conducted calculation for REE enrichment in clinopyroxenes by using 1-D steady state modeling, which duplicates simple fractional melting process and influx melting process. The results indicate that LREE-enriched melts (E-MORB-like) and LREE-depleted MORB melts (N-MORB-like) were involved in formation of the present-day concordant dunite bands within the Oman mantle with various ratios of LREE-enriched melt/LREE-depleted melt; LREE-enriched melt/LREE-depleted melt ratios are high at the segment center, and they are low at the segment end. The primitive MORB melts have possibly changed to MORB through interaction with peridotites en route to the uppermost mantle, however the interaction degrees between the segment center and the segment end were different. The difference was caused by variation of temperature profile through the Mid-ocean ridge.

Keywords: Concordant dunite band, Oman ophiolite, MORB, melt/rock interaction