Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



SSS01-13 Room:302

Time:May 19 15:30-15:45

A Slump in the Trench: Tracking the impact of the 2011 Tohoku-Oki earthquake

Michael Strasser^{1*}, Martin Kolling², Christian dos Santos Ferreira², Hiske Fink², Toshiya Fujiwara³, Susann Henkel⁴, Ken Ikehara⁵, Toshiya Kanamatsu³, Kiichiro Kawamura⁶, Shuichi Kodaira³, Miriam Romer², Gerold Wefer², Science Party⁷, Science Party⁸

¹Geological Institute, ETH Zurich, Zurich, Switzerland,, , ³Japan Agency of Marine Science and Technology (JAMSTEC), Yokosuka, Japan, ⁴Geological Institute, University of Cologne, Koln, Germany, ⁵Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology (AIST), ⁶Department of Geosphere Science, Yamaguchi University, Yamaguchi City, Japan, ⁷R/V Sonne Cruise SO219A, ⁸JAMSTEC Cruise MR12-E01

Subduction earthquakes can rupture the seafloor and trigger submarine slumps, which are potentially tsunamigenic and may affect the structural evolution of convergent plate boundaries. Bathymetric and seismic data in the Japan Trench, obtained before and after the 2011, Magnitude 9 Tohoku-oki earthquake off Japan, document significant seafloor- and subseafloor changes as tectonic and geomorphic expressions of this mega-earthquake. One challenge groundtruthing geophysical data and models, however, is to access sea- and subseafloor samples, which can be dated to assign the hypothesized process to a given earthquake. Here we present multibeam bathymetry, reflection seismic, sediment core and pore-water geochemistry data from the Japan Trench, collected after the 2011 Tohoku-Oki earthquake. Analyzing the diffusive re-equilibration of disturbed SO42- profiles over time allows us to constrain that the observed sediment disturbance and mass movements, which link to bathymetric and structural changes observed in geophysical datasets, were indeed triggered by the 2011 earthquake. We conclude that a slump in the trench significantly impacted the geometry and evolution of the shallow plate boundary system by emerging a submarine fold-and-thrust belt and abruptly narrowing the trench up to 3 km in width.

Keywords: Japan Trench, Submarine Mass Movements, Sediment core, Sulphate, R/V Sonne Cruise SO219A, RV Mirai Cruise MR12-E01