Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



SSS24-08

会場:303

時間:5月20日11:30-11:45

東北地方太平洋沖で発生した地震からの反射 sP 波の振幅分布 Amplitude distribution of sP reflected phase from offshore earthquakes in the Pacific side of Tohoku district

小菅 正裕 ^{1*} Masahiro Kosuga^{1*}

1 弘前大学理工学研究科

We observed a prominent phase on vertical seismograms of interplate and intraplate earthquakes offshore Miyagi prefecture with depth ranges from 30 to 60 km. The phase that appears between P- and S-waves is observed widely at stations in the Japan Sea side. Here we examine the waveform and travel times of this phase using Hi-net data and estimate its origin, and discuss potential usage of the phase.

A polarization analysis indicates that the phase has a strike toward the epicenter, nearly vertical dip angle, and large rectilinearity. This indicates that the phase is P-wave coming from the direction of the source. The travel time of the X-phase is proportional to epicentral distance with an apparent velocity of about 7 km/s. This suggests that the reflection/conversion occurs at relatively shallow part. There is no significant azimuthal variation in arrival times of the X-phase, which implies that the plane of reflection/conversion is nearly horizontal. We estimate the position of conversion/reflection by using ordinary hypocenter location method assuming that the phase is P-wave from the conversion/reflection point. The location is near the surface of the coastal area of Miyagi prefecture. The above observational facts of large amplitude, polarization characteristics, apparent velocity, azimuthal variation of arrival times, and the location of conversion/reflection point, are all consistent with an interpretation that the phase is sP reflected phase from the surface. This phase has already found by previous studies and has been used as a depth phase to improve the depth accuracy in hypocenter location and delineate a seismicity pattern along the plate boundary.

Next we investigated amplitude distribution of sP phase. We measured amplitude on RMS envelope as the deviation of smoothly varying envelope. In many cases the amplitude is largest at stations in Akita and Yamagata prefectures, while the amplitude is smaller at stations in the northern and southern part of Tohoku district, and at stations in the Pacific side. The focal mechanisms of these events are reverse faulting with N-S strike. In the case of reverse faulting earthquake with E-W strike, the area of large amplitude shifts to the northern part of Tohoku district. This indicates that the amplitude distribution of sP phase depends on focal mechanisms. Thus the amplitude of sP phase has a potential usage to determine focal mechanisms of offshore earthquakes, which is difficult from the P-wave polarization only.

Acknowledgement: We thank the National Research Institute for Earth Science and Disaster Prevention (NIED) for providing waveform data from Hi-net.

キーワード: sP波, 反射, 振幅, 震源メカニズム

Keywords: sP wave, reflection, amplitude, focal mechanisms

¹Graduate School of Sci. and Tech., Hirosaki Univ.