Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



STT55-04

時間:5月22日10:00-10:15

2011 年噴火後の新燃岳における無人ヘリ繰り返し空中磁気測量 Repeated aeromagnetic survey of Shinmoe-dake, Kirishima volcano, Japan, after the 2011 eruption using unmanned autonomo

小山 崇夫^{1*}, 金子 隆之¹, 大湊 隆雄¹, 柳澤 孝寿², 渡邊 篤志¹, 武尾 実¹ Takao Koyama^{1*}, Takayuki Kaneko¹, Takao Ohminato¹, Takatoshi Yanagisawa², Atsushi Watanabe¹, Minoru Takeo¹

¹ 東京大学地震研究所,²海洋研究開発機構 地球内部ダイナミクス領域

¹Earthquake Reserach Institute, Univ. Tokyo, ²Institute for Research on Earth Evolution, Japan Agency for Marine-Earth Science and Technology

We conducted the highly resolved aeromagnetic surveys around Shinmoe-dake, Kirishima volcano twice by using the unmanned autonomous helicopter, when it was after the magmatic sub-plinian eruptions.

First survey was carried out in the end of May covering 3 by 3 km area of western Shinmoe-dake. The flight altitude is as low as about 100 m above the ground and spacing of flight lines are as close as almost 100 m. Total flight distance is about 85 km. The cesium optically-pumping magnetometer was installed by hanging below the helicopter to measure the geomagnetic total intensity. Due to this survey, we detected a large geomagnetic total intensity anomaly as 1000 nT. Using these data, the horizontal map of magnetization intensity was obtained beneath the survey area. We found the followings;

1) The average value of the magnetization is as low as 1.5 A/m;

2) Northwestern Shinmoe-dake has lower magnetization than average;

3) Strong magnetization appears the area corresponding the lava at the foot of Karakuni-dake.

Second survey was carried out about 5 months later than the first survey, in the end of October to the beginning of November, in order to detect temporal change of geomagnetic total intensity. By comparing both data, we could detected a dipole-like change as large as about \pm 100nT around the crater of Shinmoe-dake. This change can explain by magnetization enhancement with 2.3 x 10⁷ Am² in the crater. In the 2011 eruptions of Shinmoe-dake, a large pancake-like lava was found in crater with 1.5 x 10⁷ m³ (Nakada et al., submitted), and the cooling of this lava probably causes this magnetization enhancement.

キーワード:繰り返し空中磁気測量,無人ヘリコプター,新燃岳

Keywords: repeated aeromagnetic survey, unmanned autonomous helicopter, Shinmoedake