

## Hayabusa-2 sampler: Surface variety of near-Earth C-type asteroid

Shogo Tachibana<sup>1\*</sup>, Hiroataka Sawada<sup>2</sup>, Ryuji Okazaki<sup>3</sup>, Yoshinori Takano<sup>4</sup>, Chisato Okamoto<sup>2</sup>, Hajime Yano<sup>2</sup>, Kosuke Kurosawa<sup>2</sup>, Koji Wada<sup>5</sup>, Hayabusa-2 sampler team<sup>2</sup>

<sup>1</sup>Department of Natural History Sciences, Hokkaido University, <sup>2</sup>Japan Aerospace Exploration Agency, <sup>3</sup>Department of Earth and Planetary Sciences, Kyushu University, <sup>4</sup>Japan Agency for Marine-Earth Science and Technology (JAMSTEC), <sup>5</sup>Planetary Exploration Research Center, Chiba Institute of Technology

Samples from C-type asteroids, which are abundantly present in the asteroid belt and of which reflectance spectra resemble those of carbonaceous chondrites, may well preserve the information covering the long history of the solar system, compared to other bodies such as comets and Itokawa-like bodies. Their scientific values will be significantly increased for return samples obtained with detailed geological contexts. Moreover, surface samples from near-Earth C-type asteroids will provide insights into the space weathering of C-type asteroids and the surface thermal processes due to irradiation of sunlight, which cannot be obtained from meteorites and interplanetary dust particles. The Hayabusa-2 is a sample return mission from a near-Earth C-type asteroid 1999JU3 (2014-2020). Here we describe a sampling system of the Hayabusa-2 spacecraft to obtain samples from multiple surface locations of the asteroid with minimal contamination and a possible sampling strategy.

Keywords: Hayabusa-2, asteroid, sample return, space exploration