

Possible interactions between Indian Ocean Dipole and intraseasonal variability in the tropical Indian Ocean

Yukio Masumoto^{1*}

¹JAMSTEC

Variations in the tropical Indian Ocean cover a wide range of spatial and temporal scales. Indian Ocean Dipole (IOD) is one of the dominant interannual climate modes there, and several intraseasonal variations are believed to have interaction with IOD through oceanic bridges and/or air-sea interactions. Two of such examples will be explored in this presentation.

The first example is influence of meso-scale ocean eddy activity in the southeastern tropical Indian Ocean, which is generated by baroclinic instability in the northern part of the South Equatorial Current. Strong meridional temperature gradient associated with the IOD event results in anomalously energetic eddy activity. This causes stronger-than-normal northward eddy heat transport, which tends to reduce the meridional temperature gradient, hence a negative feedback on the IOD event itself.

Another example is impact of intraseasonal equatorial Kelvin waves on initiation of the IOD events, particularly in 2006 case. During May to August, before the 2006 IOD event, several upwelling equatorial Kelvin waves were excited by easterly wind anomalies in the equatorial central Indian Ocean. Negative subsurface temperature anomalies at the thermocline depth appeared associated with penetration of these Kelvin waves along the Sumatra coast, creating favorable conditions for cooling of the surface layer due to monsoonal upwelling in August. Constructive interaction between the two processes may set a critical condition for the generation of the 2006 IOD.

Such scale-interactions in the tropical Indian Ocean should be investigated in more detail for better understanding of mechanisms responsible for the IOD evolution and the skillful prediction of IOD.

Keywords: Indian Ocean Dipole, Intraseasonal variations, interactions, meso-scale eddy, Kelvin waves

Influence of the Wyrтки Jets on the western Arabian Sea upwelling region

Tomoki Tozuka^{1*}, Motoki Nagura², Toshio Yamagata²

¹Graduate School of Science, The University of Tokyo, ²Japan Agency for Marine-Earth Science and Technology

The sea surface temperature (SST) in the upwelling region along the western boundary of the Arabian Sea is known to influence the Indian summer monsoon rainfall. In this study, we examine how a reflection of semiannual Kelvin waves, which is forced by westerly winds during monsoon breaks and accompanied by the Yoshida-Wyrтки Jet, may influence this region based on ocean general circulation model experiments. When results from two experiments with and without a damping near the eastern equatorial Indian Ocean are compared, the SST in the western Arabian Sea becomes colder by as much as 0.4 degree C in the latter experiment. By calculating mixed layer heat balance, it is shown that this SST difference is mainly due to a difference in horizontal advection, but is damped by surface heat flux.

Keywords: Tropical Indian Ocean, Ocean general circulation model, Equatorial wave

Role of Diurnal Warm Layers in the Diurnal Cycle of Convection over the Tropical Indian Ocean during MISMO

Hugo Bellenger^{1*}, Kunio Yoneyama¹, Yukari Takayabu², Tomoki Ushiyama³

¹RIGC - JAMSTEC, ²AORI - University of Tokyo, ³ICHARM - PWRI

The role of air sea interaction in the diurnal variations of convective activity during the suppressed and developing stages of an intraseasonal convective event is analyzed using in situ observations from the Mirai Indian Ocean cruise for the Study of the Madden Julian oscillation (MJO)-convection Onset (MISMO) experiment. For the whole period, convection shows a clear average diurnal cycle with a primary maximum in the early morning and a secondary one in the afternoon. Episodes of large diurnal sea surface temperature (SST) variations are observed because of diurnal warm layer (DWL) formation. When no DWL is observed, convection exhibits a diurnal cycle characterized by a maximum in the early morning, whereas when DWL forms, convection increases around noon and peaks in the afternoon. Boundary layer processes are found to control the diurnal evolution of convection. In particular, when DWL forms, the change in surface heat fluxes can explain the decrease of convective inhibition and the intensification of the convection during the early afternoon.

Keywords: Diurnal Warm Layers, Madden-Julian Oscillation, Preconditioning, MISMO, convection

Cluster analysis of the intraseasonal convection and its impact on the tropical tropopause temperature

Eriko Nishimoto^{1*}, Masato Shiotani²

¹Graduate school of science, Kyoto University, ²Research Institute for Sustainable Humanosphere, Kyoto University

This study investigates space-time variations of the tropical convective activities and temperatures around the tropical tropopause associated with the intraseasonal oscillation (ISO) during the southern summer by using outgoing longwave radiation (OLR) data from the National Oceanic and Atmospheric Administration and atmospheric fields from the European Centre for Medium-range Weather Forecasts Interim reanalysis data.

Cluster analysis is conducted in order to classify ISO types according to both the phase speed and the longitudinal extent of the eastward propagation. In performing cluster analysis we use the locus of convective activities observed in the unfiltered OLR data by retaining both the ISO and seasonal mean components to investigate interaction between the two. Then, the 72 ISO events in the 32 southern summers are mainly grouped into four clusters. Two of the clusters exhibit the slow (<2 m/s) propagation speed in the unfiltered OLR field and the others fast (~4 m/s). One cluster characterized as the El Nino phase has the fast speed while passing over the date line and another as the La Nina phase has the slow speed while propagating to ~120E. Compared with the other two clusters characterized as the weak El Nino-Southern Oscillation phase, the speed is slow while propagating to ~135E when the SSTs over the Western Pacific are relatively low.

Low temperatures around the tropical tropopause appear to the east of the eastward-propagating convection in the tropics and to the west in the subtropics, forming a horseshoe-shaped structure. The strength of the horseshoe-shaped temperature structure is determined by that of the convective activities. Furthermore, the strength and location of the 100-hPa temperature minima differ among the clusters. This study implies that the different ISOs would cause different impacts on the dehydration process in the tropical tropopause layer depending on their types.

Keywords: Intraseasonal Oscillation, Madden-Julian Oscillation, Cluster Analysis, Teleconnection, Tropical Tropopause Layer

Synchronization of thunderstorm activities and OLR in tropical regions

Yukihiro Takahashi^{1*}, Yusuke Sanmiya¹, Mitsuteru Sato¹

¹Dept. CosmoSciences, Hokkaido University

Based on Global ELF observation Network (GEON) and Outgoing Longwave Radiation (OLR) intensity, we carried out a correlated analysis between the number of the lightning strokes and cloud variation in the tropical regions, focusing the variation around one month periodicity. It was found that the number of lightning strokes in the Maritime Continent (MC) varies with about month periodicity in the period from February to June 2004 and shows positive correlation ($R=0.8$) with OLR in the Western Pacific Warm Pool (WPWP). That is, when thunderstorm activity in the MC is enhanced, the OLR in WPWP becomes large, meaning less cloud amount. On the other hand, OLR in the central Africa shows negative correlation with the number of lightning strokes in the MC in that period ($R=-0.7$). Furthermore, in the central Africa OLR seems to reflect the number of lightning strokes, showing good correlation between them. This implies that the activities of thunderstorms both in the central Africa and in the MC oscillate in the same phase. Such a synchronization of thunderstorms or cloud amount in global scale without phase difference has not been reported and seems difficult to explain these phenomena by conventional theories. We may need to consider the variation of solar activity, such as UV or galactic cosmic rays, whose variation in the present period (Feb-Jun 2004) shows good correlation with OLR variations in tropical region.

Keywords: thunderstorm, OLR, tropical region, synchronization

Relationships among Lightning, Precipitation, and Hydrometeor Characteristics in Equatorial Indonesia

Marzuki Marzuki^{1*}, Hiroyuki Hashiguchi¹, Masayuki Yamamoto¹, Shuichi Mori², Manabu D. Yamanaka², Yukihiro Takahashi³

¹Research Institute for Sustainable Humanosphere, Kyoto University, ²Japan Agency for Marine-Earth Science and Technology (JAMSTEC), ³Hokkaido University, ⁴Department of Physics, Andalas University, Indonesia

1. Introduction

Knowledge of the lightning activity is an important tool to the meteorologists. Many literatures have discussed on lightning activity for different regions. However, the studies pertaining to the variability of lightning occurrences in the equatorial Indonesia are scanty, except a few studies. Hence, in the present study, the variability of lightning activity in the equatorial Indonesia is examined, particularly at Kototabang (KT; 100.32E, 0.20S), Pontianak (PT; 109.37E, 0.00S), Manado (MN; 124.92E, 1.55N) and Biak (BK; 136.10E, 1.18S).

2 Data and Methodology

Lightning activity are observed from the World Wide Lightning Location Network (WWLLN) data. Following the advice of the WWLLN developers, only those lightning locations that triggered at least five sensors and that had residuals < 30 ms are included in this analysis [1]. The surface precipitation and the profiles of hydrometeors and latent heating are obtained from the products of the Tropical Rainfall Measuring Mission (TRMM) satellite. The aerosol data are retrieved from the Moderate Resolution Imaging Spectroradiometer (MODIS). Raindrop size distribution (DSD) at the surface is from a network of Parsivel disdrometers. 1.3 GHz wind profiler data at the four sites are used to determine the precipitating cloud type.


3 Results

The evidence of regional variation of precipitation microstructure (e.g., DSD) is clearly observed, and become more obvious during heavy rain. The composite spectra of PT and KT were much broader than Marshall-Palmer distribution, in contrast to the DSD at MN and BK, where the DSDs were narrow. This characteristic is consistent with the lightning activity. Figure shows regional variability of percentage occurrence of flashes. The convective storms are more intense at PT than other three sites and produce larger raindrops. Detailed analysis regarding the relationships among lightning, precipitation, and hydrometeor characteristics in equatorial Indonesia will be presented in the meeting.

References

[1] Abarca, S. F., Corbosiero, K. L., and Galarneau Jr., T. J.: An evaluation of the Worldwide Lightning Location Network (WWLLN) using the National Lightning Detection Network (NLDN) as ground truth, *J. Geophys. Res.*, 115, D18206, doi:10.1029/2009JD013411, 2010.

Keywords: Equator, Lightning, Precipitation



Origin of seasonal predictability for summer climate over the Northwestern Pacific

Yu Kosaka^{1*}, XIE, Shang-Ping¹, LAU, Ngar-Cheung², VECCHI, Gabriel A.²

¹Scripps Institution of Oceanography, ²GFDL, NOAA

Summer climate in the Northwestern Pacific (NWP) displays large year-to-year variability, affecting densely populated Southeast and East Asia by impacting precipitation, temperature and tropical cyclones. The Pacific-Japan (PJ) teleconnection pattern provides a crucial link from the tropics of high predictability to East Asia. Using coupled climate model experiments, we show that the PJ pattern is the atmospheric manifestation of an air-sea coupled mode spanning the Indo-NWP warm pool. In this coupled mode, the PJ pattern forces the Indian Ocean (IO) via a westward propagating atmospheric Rossby wave. In response, IO sea surface temperature (SST) feeds back and reinforces the PJ pattern via a tropospheric Kelvin wave. Ocean coupling increases both the amplitude and temporal persistence of the PJ pattern. Cross-correlation of ocean-atmospheric anomalies confirms the coupled nature of this PJIO mode. El Nino-Southern Oscillation (ENSO) is a major external driver of the PJIO mode, leaving the last echoes of ENSO in the IO-NWP in the form of this mode. We further demonstrate that the PJIO mode is indeed highly predictable, giving hopes for skillful seasonal forecast over the densely populated region.

Keywords: air-sea coupled mode, climate variability, East Asian summer monsoon, El Nino-Southern Oscillation

Recent progress in the MIROC5 seasonal prediction system and predictability of two flavors of El Nino

Yukiko Imada^{1*}

¹Atmosphere and Ocean Research Institute, the University of Tokyo

This study investigates the difference of the seasonal predictability for two prominent types of El Nino, traditional eastern Pacific (EP) events and central Pacific (CP) events.

We developed a seasonal prediction system using the coupled atmosphere-ocean general circulation model (AOGCM) MIROC5 co-developed by Atmosphere and Ocean Research Institute (AORI), National Institute for Environmental Studies (NIES), and Japan Agency for Marine-Earth Science and Technology (JAMSTEC). The spatial resolution is a horizontal triangular spectral truncation at total wave number 85 (T85) with 40 vertical layers, and eight ensemble forecast members are generated according to the protocol of the WCRP Climate-system Historical Forecast Project (CHFP). Hindcast products for the period 1979-2011 show high predictability of tropical climate signals with the significant anomaly correlation coefficient skill scores, even though the ocean anomaly data assimilation is applied to the initialization process. The monsoon and Indian Ocean Dipole (IOD) indices also show predictable signals until a few months later. Interestingly, our seasonal prediction system is less affected by the "spring prediction barrier" compared to most of the other AOGCMs.

We assess the difference of the seasonal predictability for two prominent types of El Nino, traditional EP events and recent CP events. Overall, the predictable months of CP events are shorter than EP events because CP events have less amplitude and are sensitive to atmospheric noises. It seems that this difference in predictability connects to the recent low predictability after 2000 as shown in Barnston et al. (2012). Characteristics of each error-growing process are also investigated.

Barnston, A. G., M. K. Tippett, L. L'Heureux, S. Li, and D. G. DeWitt (2012), Skill of real-time seasonal ENSO model predictions during 2002-11, *BAMS*, 631-651.

Keywords: seasonal prediction, ENSO, AOGCM, predictability

Study on a relationship between New Guinea coastal upwelling in the Bismarck Sea and onset of El Nino events

Takuya Hasegawa^{1*}, Toru Miyama², Jing-Jia Luo³, Bunmei Taguchi⁴, Ayako Seiki²

¹JAMSTEC/RIGC, ESC, ²JAMSTEC/RIGC, ³Bureau of Meteorology/Australian Government, Australia, ⁴JAMSTEC/ESC

We investigate oceanic and atmospheric variations related to coastal upwelling along New Guinea Island north coast before onset of El Nino event. In our previous studies, existence of the coastal upwelling and related SST cooling near New Guinea Island during December 2001 to January 2002, which was prior to onset of 2002/03 El Nino event, were shown by observational data. Furthermore, we explored cooling mechanism related to the coastal upwelling using high-resolution OGCM (OFES) hindcasts using NCEP/NCAR forcing and QuikSCAT forcing during 1981-2010. In this study, we analyze long-term output of 200-year simulation of high-resolution air-sea coupled general circulation model (SINTEX-F ver2). SST patterns similar to the observed coastal upwelling along north coast of New Guinea Island appear before onset of El Nino-like events in the SINTEX-F ver2 simulation. At those periods, positive zonal SST gradients in the western equatorial Pacific also appear in association with the SST cooling near New Guinea Island. Relatively strong westerly surface winds, which are expected by the positive SST zonal gradients, also appear in those periods. Such oceanic and atmospheric relationship is similar to that of observation for SST cooling period of December 2001 to January 2002. It might suggest that the SST cooling along north coast of New Guinea Island can relate to El Nino onset via atmospheric changes. We will also explore upper-ocean structure during the SST cooling period to further explore a relationship between the SST cooling pattern and New Guinea Island coastal upwelling before El Nino onsets using the SINTEX-F ver2 simulation.

Keywords: Air-sea interaction, coastal upwelling, Pacific warm water pool, El Nino onset

Regional Climate Modeling Study of Wind Variations over Western Pacific Warm Pool before El Nino Onsets

Toru Miyama^{1*}, Takuya Hasegawa²

¹JAMSTEC RIGC/APL, ²JAMSTEC RIGC/ESC

Regional climate modeling is an effective way to study on phenomena which found interesting in global GCMs. Regional models can play a complementary role to global models in experimental designs. While global models are free running, but may suffer from biases, regional models are bounded at lateral boundaries and achieve better realism.

Using the data of observations and global models, Hasegawa et al (2009, 2010, 2011) found strong northwesterly surface winds and cold-water upwellings along the northern coast of New Guinea often occur before El Nino onsets. They hypothesized that the cold waters generate positive zonal sea surface temperature (SST) gradient together with high SST east of the warm pool in the Western Pacific Ocean contribute to enhancement of the westerly surface winds, leading to onsets of El Nino events. The goal is to understand this region in an ocean-atmosphere coupled system. As a first step, we have conducted experiments with a regional atmospheric model. The model used in this study is the International Pacific Research Center (IPRC) Regional Atmospheric Model (iRAM) to understand the effects of the cold SST. The model covers the western Pacific Ocean with a horizontal resolution of 0.25 degree. We particularly focused on December 2001 to January 2002, as Hasegawa et al. (2009) did in their diagnostic study. The model well reproduced events of wind westerly surface winds in this region. Experiments show that wind variations near the New Guinea are responsive to local SST. Even when the lateral boundary condition is unchanged, westerly surface wind is weakened when the cold signal by the upwelling is eliminated from the SST field. We also pay attention to the role of the high mountains of New Guinea in shaping climate around this region. An experiment showed the orography of New Guinea causes rising air motion above the mountains. Recent experiments of the coupled ocean-atmospheric model (coupled to the HYbrid Coordinate Ocean Model using the Earth System Modeling Framework) will be also reported.

Keywords: El Nino, Pacific Ocean, regional model, upwelling, Tropics, convection

Off-equatorial influences on equatorial Atlantic variability

Ingo Richter^{1*}, Swadhin Behera¹, Yukio Masumoto¹, Bunmei Taguchi², Hideharu Sasaki², Toshio Yamagata³

¹Research Institute for Global Change, JAMSTEC, ²Earth Simulator Center, JAMSTEC, ³Application Laboratory, JAMSTEC

Interannual-to-decadal variability in the tropical Atlantic is dominated by two modes of variability. The zonal mode governs in the equatorial Atlantic and is thought to rely on dynamics akin to El Niño-Southern Oscillation (ENSO). The meridional mode, on the other hand, involves sea-surface temperature (SST) anomalies in the northern and southern tropical Atlantic centered at 15°N and 15°S, respectively.

In the present study we use the NCEP reanalysis, OFES hindcast, and CSIRO Mk 3.5 coupled GCM to reexamine the dynamics governing the zonal mode. We find that equatorial wind stress forcing and ENSO-like dynamics can explain some of the observed warm events but not all of them. In particular there are warm events that occur despite easterly surface wind anomalies in the preceding months. This is due to sub-surface warm waters being advected from approximately 5°N toward the equator. The sub-surface warming is ultimately related to SST anomalies in the northern tropical Atlantic, which induce wind stress curl anomalies that force downwelling just north of the equator. This suggests a mechanism by which off-equatorial ocean conditions can influence the zonal mode of variability and poses an additional challenge to skillful predictions in the region.

Keywords: tropical Atlantic, equatorial Atlantic, zonal mode, meridional mode, meridional advection

Locally amplified Ningaloo Nino off the western coast of Australia

Takahito Kataoka^{1*}, Tomoki Tozuka¹, Swadhin Behera², Toshio Yamagata²

¹The University of Tokyo, ²JAMSTEC

Using observational and reanalysis data, the mechanism of a new climate mode off western Australia called "Ningaloo Nino" is investigated. It is associated with positive sea surface temperature (SST) anomalies and peaks during austral summer. There are two types of Ningaloo Nino: The "locally amplified" and "non-locally amplified" events. The former can develop through an intrinsic unstable air-sea interaction off western Australia; an anomalous cyclone generated by positive SST anomalies forces northerly alongshore wind anomalies, which induce coastal downwelling anomalies, and enhance the warm SST anomalies.

It is found that the locally amplified Ningaloo Nino cause positive rainfall anomalies along the coast of western Australia, but the signals are subtle and the northern part tends to become drier because of a weaker monsoon.

Keywords: Ningaloo Nino, unstable air-sea interaction, coastal upwelling, western Australia, precipitation

ENSO simulation in GCMs: A review and recent progress

Masahiro Watanabe^{1*}

¹AORI, The University of Tokyo

Changes in the activity of the El Niño-Southern Oscillation (ENSO) phenomenon under global warming could potentially have a large impact on the global weather, society, and economy, but have not so far been converged in ensembles of future scenario experiments based on multiple climate models. Using parameter ensembles of four high-end climate models with smaller errors in simulating ENSO in the present climate, we demonstrate that the ENSO will become more energetic under global warming. This occurs because the mean atmospheric state in a warmed climate is wetter over the central-eastern equatorial Pacific. The wetter mean state favours an eastward shift in the equatorial zonal wind stress response to El Niño/La Niña, which acts to increase the ENSO amplitude due to enhanced coupled instability. A careful analysis of the previous multi-model ensemble suggests a similar mechanism at work and indicates that the precipitation increase over the cold tongue region, relative to the change over the entire equatorial Pacific, is the key factor for the robust intensification of ENSO. A preliminary analysis to the CMIP5 archive, however, does not necessarily support the above conclusions.

Keywords: GCM, ENSO, Global warming

Mechanism for the asymmetry in ENSO transition and duration

Masamichi Ohba^{1*}

¹Central Research Institute of Electric Power Industry (CRIEPI), Environmental Science Research Lab.

The El Nino-Southern Oscillation (ENSO), which consists of a quasi-periodic (3-7-yr timescale) warming (El Nino) and cooling (La Nina) of the tropical central and eastern Pacific Ocean (CEP), is the most dominant driver for Earth's interannual climate variability. ENSO prediction is of practical interests, in addition to scientific, because of its large environmental and societal impacts. To predict and understand the variability of ENSO, a number of investigators have suggested conceptual theories, providing a comprehensive idea regarding the cyclic nature of ENSO. The mechanisms in these conceptual theories effectively capture the observed phase transition from El Nino to La Nina and successfully reproduce the linear oscillation of ENSO. However, several studies have reported that a type of break in the ENSO cycle occurs when La Nina shifts to El Nino. The air-sea coupled system over the Pacific somehow remains in a weak La Nina state for up to two years, while El Nino tends to turn rapidly into La Nina after the mature phase. Recent studies (Ohba and Ueda 2009; Ohba et al. 2010; Okumura et al. 2011) have reported that the nonlinear atmospheric response to the CEP sea surface temperature forcing is a fundamental cause of the asymmetry in the transition. Because the duration of an ENSO episode can cause severe drought, for example, the 1999-2001 drought in central Asia from, and this duration is difficult to reproduce in most coupled general circulation models, understanding of the ENSO asymmetry is important for improving seasonal climate forecast skills (Ohba et al. 2010; Ohba and Watanabe 2012). Therefore, the asymmetry of transition/duration is an important aspect of ENSO. In the presentation, these recent studies will be introduced with the discussion of its long-term change and asymmetry in the ENSO predictability.

Keywords: El Nino/Southern Oscillation, Pacific Ocean, Indian Ocean

Prediction and Projection of Tropical Cyclone Activity over the Western North Pacific Using CMIP5 Near-Term Experiments

Masato Mori^{1*}, KIMOTO, Masahide¹, ISHII, Masayoshi², WATANABE, Masahiro¹, MOCHIZUKI, Takashi³

¹Atmosphere and Ocean Research Institute, University of Tokyo, ²Meteorological Research Institute, ³Japan Agency for Marine-Earth Science and Technology

In line with the experimental design for near-term climate prediction toward the 5th Assessment Report of the Intergovernmental Panel on Climate Change, we performed ensembles of initialized decadal hindcast and near-future (NF) projection using three versions of the coupled atmosphere-ocean model MIROC. In this study, interannual and multiyear predictability of tropical cyclone (TC) activity in the western North Pacific (WNP) is explored, using the initialized hindcasts. In addition, global warming impacts on WNP TC activity in the NF are also examined using the NF projection up to 2035.

The hindcasts show that year-to-year variation of TC number reasonably captures the observation. Interannual variability for TC genesis and occurrence frequency (TGF and TOF) associated with El Niño Southern Oscillation (ENSO) is found to be predictable mainly through better prediction of sea surface temperature (SST) and lower-tropospheric large-scale vorticity anomalies. These results indicate that models are able to reproduce the major basic mechanisms that link TC genesis with large-scale circulation. On the multiyear timescale, skillful prediction of TC number is likely difficult at least in our hindcasts, but three-year-mean states of hindcast started in 1998 reasonably capture observed major characteristics of TC activity associated with the Pacific climate shift during the late 1990s through the initialization.

Projected NF (2016-2035) change in WNP TC genesis number shows significant reduction (approximately 14%) especially over the western WNP even in the NF when the global warming is not so prominent compared with the end of this century. The reduction is likely due to the suppression of large-scale lower-tropospheric vorticity and relative humidity, and enhancement of vertical wind shear. The projected SST exhibits a more pronounced warming over the eastern tropical Pacific and accompanies weakening of Walker circulation via redistribution of tropical convection activity, which appears to be responsible for the change in large-scale fields in WNP.

Keywords: Tropical Cyclone, Predictability

Slowdown of the Walker circulation driven by tropical Indo-Pacific warming

Hiroki Tokinaga^{1*}, Shang-Ping Xie², Clara Deser³, Yu Kosaka², Yuko M. Okumura⁴

¹IPRC, University of Hawaii, ²Scripps Institution of Oceanography, University of California at San Diego, ³National Center for Atmospheric Research, ⁴Institute for Geophysics, The University of Texas at Austin

A suite of ship observations including sea level pressure, marine cloud, surface wind, and ocean subsurface temperature show that the Walker circulation has slowed down for the past century. The cause of this slowdown is investigated using a multi-model ensemble of atmospheric GCM simulations forced by several datasets of historical SST. The models reproduce observed changes well if the right SST datasets are used. The results show that the Walker circulation change over the past six decades was induced mostly by changes in zonal SST gradient across the Indo-Pacific Oceans, and that the warming over the Indo-western Pacific is not as large as previously thought. The widely-used SST datasets show intense warming over the tropical Indo-western Pacific, where uncertainty of SST warming trend is especially large. As a result, atmospheric GCMs forced by the conventional SST datasets tend to strengthen the Walker circulation, in disagreement with observations. The observed circulation change over the tropical Pacific contains large natural variability but provides a useful constraint on historical SST reconstruction.

Keywords: Walker circulation, Climate change, Tropical Pacific, Tropical Indian Ocean, Ocean-atmosphere interaction

Intercomparison of CMIP5 Ocean Model Performance for SST Variations over EEIO and its Relation to Thermocline

Ibnu Fathrio^{1*}, Yasumasa Kodama¹

¹Graduate School of Science and Technology, Hirosaki University

The motivation of this study is to evaluate SST variations of Coupled Model Intercomparison Project (CMIP5) dataset in Eastern Equatorial Indian Ocean (EEIO) by considering the influence of subsurface ocean structure. Variations of SST are studied by applying Power spectral density (PSD) analysis on SST of CMIP5 dataset and observation of SODA dataset. Some models show stronger/weaker SST variations than observation on specific time scale. Based on the strength of SST variations relative to observation, models are divided into three groups: strong model, moderate model and normal model. Normal models have SST variations close to observation, while strong and moderate models show stronger SST variations (relative to observation) on 1-2 years, 2-3 years and 3-7 years time scale.

The cause of strong SST variations on 3-7 years time scale is related to shallow thermocline of models. Strong linearity in SST-thermocline relation may indicate more dominant influence of subsurface to SST variations on this time scale. In warming climate, relationship between thermocline and SST is still maintained; models with shallow thermocline show stronger SST variations than models with deeper thermocline. Many models show unchanged thermocline depth, which may become the cause of a little change in SST variations on 3-7 years time scale.

Keywords: CMIP5, Eastern Equatorial Indian Ocean, Thermocline, SST variations

A distinct stronger warming in the tropical tropopause layer during 2000s: Association with minor volcanic eruptions

Sanjay Mehta^{1*}, Massatomo Fujiwara², Toshitaka Tsuda¹

¹Research Institute for Sustainable Humanosphere (RISH), Kyoto University, Japan, ²Faculty of Environmental Earth Science, Hokkaido University, Japan

The trends and various interannual variability components in the tropical tropopause layer (TTL) over the tropics (15oS-15oN) are examined by employing upper air data from GPS Radio Occultation (RO), radiosonde (IGRA, RICH and HadAT2) and ERA-Interim during 2001-2010. The detection capability of the GPS RO, though with limited data coverage, has been shown in previous studies. The temperature anomalies from unadjusted radiosonde (IGRA), adjusted radiosonde (RICH and HaAT2), and ERA-Interim shows favorable comparison with GPS RO except at 100 hPa in ERA-Interim data. Detail analysis of the warming observed in the TTL during 2001-2010 using both standard linear and multiple regressions is carried out. The temperature trend estimated using standard linear regression analysis (i.e. allowing the contributions from various interannual variability) reveals a strong warming of about 0.5-1.5 K/decade in the TTL (about 16-19 km) with maximum warming at about 18 km in each data during 2001-2010. Further, multiple regression analysis is performed while including various interannual components such as Quasi-Biennial Oscillation (QBO), El Nino Southern Oscillation (ENSO) and stratospheric Aerosol Optical Depth (AOD). We performed two types of multiple regression analysis considering without (method-1) and with (method-2) seasonal modulation of the interannual components. The distinct warming in the TTL is partially but not completely removed on removing the QBO and ENSO components. However, on removing the AOD along with QBO, ENSO removes the distinct warming in the TTL. Therefore, this study shows that the strong distinct warming in the TTL is associated with minor volcanic eruptions during 2000s. Positive and significant AOD responses to the temperatures of about 0.1-0.2 K/AOD-Index are observed in the TTL region which explains about 5-15% of the total variance during 2001-2010.

Keywords: Temperature trends, Climate change, Tropical tropopausue layer, El Nino Southern Oscillation, Stratospheric Aerosol Optical Depth