

Gravel and sand particle structures on bar surfaces and sediment pulse movements in the Sendai River, Tottori

Yoshinori Kodama^{1*}, Takeru Yumura²

¹Fac. Regional Sciences, Tottori Univ., ²Densyukan

A slope failure supplied a huge amount of metamorphic black gravel particles, nearly 8,000 dump trucks equivalent, to a middle part of the Sendai River in 1998. After this event, sediment pulse movements of these black particles along the Sendai River were surveyed several times. As a large flood occurred in 2011 after a long time in the Sendai River, we conducted our survey on sediment pulse movement again; detecting ratios of black particle on bar surfaces through sieving gravel according to phi scale, classifying between black particles and others, and weighing them. We also recorded gravel and sand particle sedimentary structures after Kodama (1994) on bar surfaces through a line sampling method. The result shows intermittent 13 km movements of sediment pulses of 16-32 mm & 32-64 mm particles over 13 years. Moreover, a large flood in 2011 had changed gravel and sand particle sedimentary structures from sand rich conditions to sand starved ones. Sediment pulse movements along the Sendai River depend not only on flood discharges but also on gravel particle sedimentary structures which relate to "sand rich" or "sand starved" conditions on gravel bar surfaces: that is size mixture effects.

Keywords: sediment pulse movement, gravel and sand sedimentary structure, surface gravel on river bed, size mixture effect, the Sendai River in Tottori, gravel particles of the Sangun Metamorphic rocks

Landform changes and flood condition in the upper reaches of the River Azusa, central Japan

Hiroshi Shimazu^{1*}

¹Rissho University

The upper reaches of the River Azusa in central Japan is a braided gravel-bed river running down Japan Alps. They are characterized by frequent landform changes occurring in the riverbed. This study aims to clarify geomorphic processes of the landform changes of the riverbed using geomorphological maps and images of the observation site. The maps were made by surveying in every year from 2007 to 2012. The images were taken with interval shooting cameras at intervals of 10 to 30 minutes in daytime from July to October in 2011 and from June to October in 2012. Major landform changes, such as channel migrations, occurred in 2009, 2010 and 2011. No major landform change occurred during the working periods of the interval shooting cameras. The surveying for the geomorphological maps were carried out in August and October in every year. During the period only small change of landforms was recognized. They occurred in severe heavy rain events more than 120 millimeters per day during the snowmelt flooding season in April and May and/or the rainy season in June and July. In 4 July 2011 water level became higher caused by 50 millimeters per day rainfall. Although on 20 September 2011 more than 140 millimeters per day rainfall was recorded, water level does not rise bankfull stage and only slight landform changes take place. Based on the analysis of geomorphological maps and the profile of the cross section of the riverbed, channel migrations were not caused by lateral shifting of the channels. Burying former channels and excavation of new channels caused channel migrations and/or channel pattern changes on the riverbed.

Keywords: gravel-bed river, landform change, geomorphic process, geomorphological map, River Azusa, Kamikochi

Landform evolution of the river mouth of Ichinomiya River, southern part of Kujukuri plain

Kuniyasu Mokudai^{1*}, Futoshi Nanayama², Shinzou Ooi³

¹Pro Natura Foundation Japan, ²AIST, ³Ibaraki Univ.

Landform evolution of river mouth of Ichinomiya River about recent 100 years is revealed by analysis from old pictorial map, old topographic map and aerial photography.

Keywords: river landform, bar, flood, river mouth

Evolution processes of the Oshika Gorge, in Tottori Prefecture

Takuma Watakabe^{1*}, Yuki Matsushi², MATSUZAKI, Hiroyuki³, Yoshinori Kodama⁴, SHINNOKI, Miho⁵

¹Graduate School of Regional Sciences, Tottori University, ²DPRI, Kyoto University, ³MALT, University of Tokyo, ⁴Fac. Regional Sciences, Tottori University, ⁵JA Bank Tottori

In general, nick-zone of a river retreats. But according to previous studies, Oshika Gorge in Misasa, Tottori-prefecture with dynamic equilibrium stage seems to incise vertically. We estimated evolution processes of the Oshika Gorge using in-situ Terrestrial Cosmogenic Nuclides dating (¹⁰Be and ²⁶Al) to determine dominant erosion mode; retreating or down cutting mode. The bedrock of the gorge consists of granite. We collected 12 granite samples from surface of erosional terraces; 8 samples were collected at A site. And We extracted one sample at a time at B~E points. At A site, strath terraces developed 7 terraces and present streambed. Incision rate of A site is calculated to be about 0.25mm/yr. But B, C and D sites are calculated to be 1.31mm/yr, 1.38mm/yr, 5.28mm/yr, respectively. E site is calculated to be 5.28mm/yr. These results suggest down cutting mode fits the Oshika Gorge from incision rate of B, C, and D. Uplifting rate of the Chugoku-Mountains estimates 0.1mm/yr (Ohmori, 1980). Incision rate of the Gorge is ten times faster than uplifting rate.

Keywords: Nick zone, Downcutting process, in-situ cosmogenic radionuclides, Incision rate, Oshika Gorge, Dynamic equilibrium

Late Quaternary development of terraces in the Kumkol Basin at the northeastern Tibet as constrained by in situ CRNs

Yoshiki Shirahama^{1*}, Yosuke Miyairi², Honglin HE³, Bihong Fu⁴, Ken-ichi Kano⁵, Tomoo Echigo⁶, Yusuke Yokoyama², Yasutaka Ikeda¹

¹Earth & Planetary Science, The University of Tokyo, ²Atmosphere and Ocean Research Institute, The University of Tokyo, ³Institute of Geology, China Earthquake Administration, ⁴Institute of Geology and Geophysics, Chinese Academy of Sciences (IGGCAS), ⁵Faculty of Science, Shizuoka University, ⁶Geo-Research Institute

In the Kumkol Basin at the northeastern margin of the Tibetan Plateau are well developed fluvial and fluvio-glacial fans and terraces, which have been deformed by many faults and folds in association with the lateral growth of the Plateau. Therefore the Kumkol Basin is one of the most important areas to understand the ongoing tectonics of the Tibetan Plateau, but has rarely been studied because of difficult accessibility. Our work is the first attempt to reveal the geomorphological development of the Kumkol Basin.

Our first field survey was carried out about fluvial and fluvio-glacial terraces (Sijiquan terraces) along the Sijiquan River, where is easier to access than other areas in Kumkol Basin. In this presentation we report the results of geomorphological analysis of the Sijiquan terraces and analyses of in situ cosmogenic radionuclides in terrace deposits, and then we discussed about the development of fluvial and fluvio-glacial landforms of this area in late Quaternary time.

Our satellite-image interpretation of the Sijiquan area revealed that terraces are classified into H, L1 and L2 groups in descending order. Morphological characteristics indicate that L1 and H groups were formed by aggradation in different periods and L2 was formed by degradation. As aggradation processes are strengthened in glacial periods because of increasing production rate of clastic materials under glacial and periglacial environments, H and L1 groups are likely to have been formed in the Penultimate-glacial and the Lastglacial periods, respectively.

We collected samples from H and L2 surfaces and present river floor (PRF) for measuring cosmogenic ¹⁰Be concentration. More than 20 quartz grains collected at the surface of each sampling location were amalgamated and then measured. The measured ¹⁰Be concentration in the PRF sample was higher than those in some samples collected from terrace surfaces, indicating that the initial value (inheritance) of ¹⁰Be concentration (i.e., the value at the time of deposition) has not been constant during the formation of the Sijiquan terraces. We estimate that present-day river gravel contains higher amount of reworked grains derived from older terraces in the drainage due to dramatic change in erosion-deposition processes. This estimation is supported by a satellite-image observation that present-day first-order gullies originate mainly from older aggradational fans within the drainage whereas the source area of aggradational fan deposits extends higher in bedrock slopes.

In order to estimate the inherited ¹⁰Be values for aggradational terrace deposits, we carried out grain-by-grain measurements of ¹⁰Be concentrations for the PRF sample; the minimum value among 15 grains can be considered as maximum value of inheritance. By using it, we calculated the minimum age of each sample, on an assumption of no surface erosion. The minimum ages of H and L2 terraces are found to be 94 ka and 15.5 ka, respectively. These ages are not in contradiction with our morphological age estimation.

In conclusion, it is likely that, in the Kumkol Basin, the fluvial regime changes from aggradational in glacial periods to degradational in interglacial periods.

Keywords: Tibetan Plateau, Qaidam Basin, Tectonic landform, Surface Exposure Dating, Fluvial terrace

Formation age and geomorphologic history of the Lonar impact crater deduced from in-situ cosmogenic Be-10

Atsunori Nakamura^{1*}, Yusuke Yokoyama¹, Yasuhito Sekine², Kazuhisa Goto³, Goro Komatsu⁴, P. Senthil Kumar⁵, Hiroyuki Matsuzaki⁶, Takafumi Matsui⁷

¹AORI, The University of Tokyo, ²Complexity Sci. & Eng., The University of Tokyo, ³International Research Institute of Disaster Science, Tohoku Univ., ⁴IRSPS, Università d' Annunzio, ⁵National Geophysical Research Institute, India, ⁶Department of Nuclear Engineering and Management, The University of Tokyo, ⁷PERC, Chiba Institute of Technology

Impact cratering is a dominant surface modification process on planetary surfaces. In the inner solar system, the large majority of impacts occur on bodies covered by primitive igneous rocks. However, most of the impacts remaining on Earth surface are on different rock types than that of the inner planet and hence geologic knowledges derived from Earth's surface cannot be translated readily. The Lonar crater is a 1.88-km-diameter crater located on the Deccan basaltic traps in India (ca. 65 Ma), and is one of the very few craters on Earth emplaced directly on basaltic lava flows. Therefore, the Lonar crater provides a rare opportunity to study impact structures observed on the basaltic surfaces of other terrestrial planets and the Moon. Since the ages of terrestrial impact structures is an key to understand geomorphological processes following to the impact, various dating methods has been applied to the Lonar Crater such as fission track (Storzer and Koeberl, 2004), radiocarbon (Maloof, 2010), thermoluminescence (Sengupta et al., 1997), and ⁴⁰Ar/³⁹Ar (Jourdan et al., 2011). Yet, a large discrepancy between these methods ranging from ca. 15 to 570 ka has been resulted. Here we report surface exposure ages based on in-situ cosmogenic ¹⁰Be in order to obtain a precise age of the Lonar crater as well as to study the geomorphologic evolution of the Lonar Crater. The samples are collected from the topographic highs on the rim of the crater and from the ejecta blanket. In-situ ¹⁰Be exposure age together with newly obtained radiocarbon age of pre-impact soil suggest potential problems of previous ages recently reported by (Jourdan et al., 2011) that ⁴⁰Ar/³⁹Ar dates are biased because of inherited ⁴⁰Ar in impact glass. Systematically young exposure age from the rim samples compared to the samples from the ejecta blanket indicate that the rim of the Lonar crater is being actively eroded. Spatial age distributions observed from the Lonar creator is not the same as the pattern reported from the well-studied Barringer crater in Arizona (Nishiizumi et al, 1991, Phillips et al., 1991), highlighting the different geomorphologic history of the two craters under different climatic and lithologic settings.

Keywords: exposure age, in-situ cosmogenic nuclide, impact crater, Lonar crater, erosion, geomorphology

InSAR-image observation of landslide surface deformation triggered by the 2008 Iwate-Miyagi Nairiku Earthquake

Hiroshi, P. Sato^{1*}, Basara Miyahara¹, Takaki Okatani¹, Mamoru Koarai¹

¹GSI of Japan

SAR (synthetic aperture radar) is a microwave imaging system used to monitor Earth's surface. We found that fringes in the image has the passibility to detect local landslide surface deformation, apart from the deformation caused directly by the earthquake-induced crustal dynamics. And the detected local deromation was considered in overlaying Kurikoma volcanic land condition map (geomorphological map) published by GSI. It was also found that the deformation area is in harmony with the area of pyroclastic deposition from Kurikoma volcano.

Keywords: earthquake, Iwate-Miyagi Nairiku earthquake, landslide, SAR

Usability of the morphometry of hummocks to estimate the volume of catastrophic sector-collapses

Hidetsugu Yoshida^{1*}

¹Meiji University

To conduct risk assessments for low-frequent but large-scale physical processes, such as catastrophic volcanic sector-collapse and associated rockslide-debris avalanches, data from similar events serve as useful reference points. This study investigates hummocky topography as a tool in the estimation of the volume of volcanic sector-collapse events, i.e., the magnitude of the collapse-to-avalanche events. The topographical differences in the debris avalanche fields that might affect the formation of hummocks are examined so as to assess the application of morphometric analysis of hummocks to estimate the volume of the sector-collapses.

Yoshida et al. (2012; *Geomorphology*) described the functional relationship between the size (plane area) and horizontal distances from the source of the hummocks on debris avalanche depositional surfaces using a regression with the equation $A = a \exp(-bD)$. Each avalanche has distinctive intercept and slope values. The intercept coefficient indicates the average initial size of hummocks at distance = 0 km, and show a high correlation with the volume of collapsed masses. Therefore, there is a high potential to calculate the initial average size of hummocks at the source area from the volume of the collapsed mass of the volcanic body. This indicates that the reverse is also possible: we can estimate the possible collapse volume solely from the hummock distributional pattern of a certain avalanche. This study assesses the applicability of the abovementioned *a*-to-volume relationship to other examples from different topographical settings.

Examples of the "wide-spread (freely spreading)" type of debris avalanches similar to those in Yoshida et al. (2012), the 1888 and the Okinajima Bandai volcano debris avalanches, were analyzed (Yoshida, 2012, 2013; TJGU). Their *a*-to-mass volume relationships are accurately described by the empirical relationship of Yoshida et al. (2012). For a similar type of avalanche, the Yotei debris avalanche, the actual *a* value is slightly larger than the *a* value calculated from the known collapse volume using Yoshida et al. (2012) equation. However, the volume of Yotei collapse may be underestimated. The author also examined a "valley-filling" debris avalanche: the Kannongawa debris avalanche. The Kannongawa debris avalanche is accurately described by the Yoshida et al. (2012) equation. Therefore, the empirical relationship proposed by Yoshida et al. (2012) can be considered applicable to many of debris avalanches, albeit with some exceptions. One of the exceptions is the Kisakata debris avalanche. The empirical relationship can be not applied to the Kisakata debris avalanche, where the hummocks are strongly regulated by the topographical complexity of the debris avalanche route.

Keywords: sector-collapse, hummocks, landforms

Threshold of uplift rate in the experiments of landform development with rainfall-erosion and uplift

Shunji Ouchi^{1*}

¹College of Science and Engineering, Chuo University


A series of experiments with rainfall-erosion and uplift of various rates suggests the existence of threshold uplift rates, across which experimental landforms show different aspects of development. A mixture of fine sand and kaolinite compacted in a square-prism-shaped container (60x60x40cm) is pushed out at a constant rate from a flat surface under the artificial mist-type rainfall. In the experiment with a low uplift rate below the lower threshold, drainage networks develop as the surface slowly uplifted. The erosion is exclusively fluvial and no slope, on which failures occur, develops. The maximum height of the surface increases with the uplift while a remnant of flat incipient surface remains, but this increase stops when the fluvial erosion becomes effective on the whole surface. The erosion on the top of low hills where the fluvial erosion works least balances with the uplift. The bottom of major valleys, where the fluvial erosion works most, is considered to have become stable at this point and no further downcut is expected. In the area between the bottom of major valleys and the top of hills, on the other hand, the erosion slightly exceeds the uplift, resulting in slow decrease of average height. A certain low characteristic relief determined by the mound erodibility and rainfall intensity develops and lasts for a long period. When the uplift rate becomes higher than the lower threshold, slopes appear at valley sides (and faults) while drainage networks are developing. After drainage networks develop, uplift exceeds erosion in the upstream area where fluvial erosion works less. Hills grow higher and slope processes, especially failures, become significant. Two types of slope failures are observed. One is relatively small failures which occur on the upper part of slopes. Surface material on the upper part of slopes absorbs much water and falls down with its own weight. Another is large scale failures observed after large and high slopes develop with rapid uplift. A whole slope collapses when its base is eroded away. The former small failures occur any time when slopes of a certain height form, and do not affect the average height unless yielded debris are carried out from the uplifted area by fluvial processes. Debris deposited in the uplifted area, however, tend to increase the gradient of streams enhancing the ability of transport. This means that sediment supply from slopes and erosion by streams would balance to form relatively stable relief and height. In the case of the latter large failures, debris from slope failures flow down as debris flows directly out from the uplifted area (even out from the deposition area when its width is narrow), and reduced relief and height significantly. This type of large slope failures occurs intensively at certain intervals depending on the rate of uplift. The surface rises and the relief increases with the uplift between the periods of intensive large slope failures, and the mound height and relief may show oscillating changes around a certain value corresponding to the rate of uplift. However, considering the magnitude of change and the length of interval, it seems difficult to call this condition 'steady state.' 'Quasi steady state' might be an adequate name for this phase of experimental landform development, including the landform development dominantly with small slope failures. If the uplift rate becomes even higher and crosses the possible upper threshold, uplift overwhelms erosion and hills grow into high mountains until the relief (and height) hits the limit determined by the width of deposition (as well as uplift) area and the strength of mound forming material. Whenever the uplift ends, the experimental landform starts decreasing its height and relief exponentially, and a peneplain-like surface appears finally after a very long period of erosion.

Keywords: rainfall-erosion experiment, landform development, threshold uplift rate, slope failures, quasi steady state, limit of mountain growth

HGM22-09

Room:101B

Time:May 21 16:45-17:00



New development of the wide soil penetrator GS-YH-13 intended to high recovery rate of excavated sample

Hideki Yoshikawa^{1*}, Futoshi Nanayama², Kiyoyuki Shigeno³

¹National Institute of Advanced Industrial Science and Technology, ²Geological Survey of Japan, AIST, ³Graduate School of Sci. and Eng., Ibaraki Univ.

In general, we use peat sampler and soil penetrator, which are common excavation methods in geomorphological investigation because of low environmental impact. Upon collecting a sample minimal excavation, although this method is a very effective. But it was difficult to get wet or/and sandy samples because of simple structure of its sampler. In addition, its weight is piling up, such as connecting rods, had been struggling to move in wetlands. Although we have been developing and improving our soil sampler over the past 10 years. We retry to improve the new wide soil penetrator GS-YH-13 in three points as bellows; (1) increasing the recovery rate of the wet and sandy samples, especially in the distal end portion which is that there is no loss of sample, (2) targeting 1m as the length of the sample while maintaining the large diameter (30mm) of sampler, (3) reducing the weight and increasing the convenience using for wetlands.

Keywords: geomorphological investigation, wide soil penetrator, GS-YH-13, high recovery rate, excavated sample, development

Internal Structure of a Lithalsa in the Akkol Valley, Russian Altai Mountains

Go Iwahana^{1*}, Kotaro FUKUI²

¹IARC / University of Alaska Fairbanks, ²Tateyama Caldera Sabo Museum

Perennial frost mounds are present near the edges of ponds located on the terrace between the mountain flank and a valley-bottom lake in the Akkol valley of the Russian Altai Mountains. These mounds are 10?50m in diameter and 3?6m in height. We describe one mound, identified as a lithalsa, which had been eroded so as to expose almost the entire vertical cross-section and reveal its internal structure. The frozen core consisted mainly of soil segments suspended in reticulate ice lenses with a mean thickness of 11?48mm and a maximum thickness of about 160mm. The shapes of the soil segments matched their neighbours. Other features included soil segments suspended in the ice veins shaped like En echelon gash veins, and the presence of a radial structure of ice-rich and sediment-rich frozen bands. These features in the core all suggest a differential stress field during and after heaving of the mound due to ice segregation, acting as a more important factor than the thermal gradient, including its direction and water supply. Based on observations of the cross-section and isotopic analysis of the ice in the permafrost core, this lithalsa appears to have developed by syngenetic differential ice segregation, even after uniform thermal and hydrologic conditions, since these were followed by epigenetic creep deformation of the cryostructure, probably with a redistribution of ice due to non-coaxial shear. Given the complexity of the cryostructures in this lithalsa, the reconstruction of its development using cores samples from boreholes would have been very difficult, particularly since the dip angles of segregated ice lenses and soil segments suspended in the ice networks have changed. Laboratory experiments and numerical simulations, supported by field observations such as those in this paper, are needed to generate further insight into the growth processes of lithalsas.

Keywords: lithalsa, frost heave, frost mound, Altai, segregated ice, internal structure

Weathering of bedrock surface after glacier retreat in the Swiss Alps

Kento Adachi¹, Atsushi Ikeda^{1*}, Tsuyoshi Hattanji¹

¹University of Tsukuba

This paper discusses how the strength of intact bedrock surface has been changed within twenty thousand years in an alpine environment. The strengths of rock surfaces were measured in the Upper Engadin, Swiss Alps, where the history of glacier retreat after the Last Glacial Maximum has been well investigated. The glacier distributions of three main stadials are evident in the study areas, which are the Little Ice Age (AD1850), Younger Dryas (12 ka) and Oldest Dryas (15-18 ka). Forty-one roches moutonnees and twenty-four rock walls of glacial troughs were selected for this research, which consist of granodiorite, diorite, gneiss or limestone. The rebound values of Schmidt hammer and Equotip hardness tester, and P-wave velocities were measured as indices of surface weathering. To compare bedrock surfaces and the intact parts, two blocks were sampled from two gneiss roches moutonnees with different ages. Their specimens, not including the weathered surface, were provided for an elastic wave test, uniaxial compression test, radial compression test and Equotip test.

These tests suggested that the strength of rock surfaces has been reduced through the last 20,000 years, except for that of limestone. The decrease in the strength is remarkable at the early stage of the weathering. In particular, the decrease rates of granodiorite and diorite can be approximated by logarithmic curves. Although granodiorite and diorite roches moutonnees exposed after the Little Ice Age are mostly unweathered, gneiss rocks with the same age have already been weathered over large parts of their surfaces. Unloading from overlying glaciers weakened the surfaces of gneiss rocks, reflecting their schistosity parallel to the surface. In contrast, the surfaces of limestone rocks have been constantly renewed by dissolution. Therefore, the weathering duration does not affect the surface strength of limestone but increases its surface roughness. The surface strength of rock walls of glacial troughs is controlled by occasional rock exfoliation owing to the gravity, as well as by weathering conditions.

Keywords: weathering, roche moutonnee, Holocene, Late Glacial, Schmidt hammer, Swiss Alps

Found Kikai-Tozuhara tephra on central district of Kaetsu-Uplands and its meaning

Tsutomu Sakamoto^{1*}

¹Kanazawa University Graduate School of Natural Science and Technology

Kaetsu uplands are located between Ishikawa and Fukui Prefecture and placed on northern part of active faults on east side of Fukui plain. This area is compressive stress field by many active faults and folds. So that this district is consist of clear upheaval and sedimentation areas. Reserching area is Central part of Kaetsu uplands. From existing research, this researching area is decided into M2 terrace. But marine terraces on east part of this area are distributed into M2 without established era data as tephra, paleomagnetism, but west part of it is decided by Youokura(1991)and so on. So, I thought that before researching all of marine terrace in this area.I have to decide whether or not all of this area is M2 terrace.

I found silt layer at Akao, Awara-city on Kaetsu uplands. Took off sands and experimented. I used MAIOT by Furusawa geology to prove an index of refraction.

I found Kikai-Tozuhara tephra(K-Tz)from these samples because

- 1) an index of refraction is between 1.396 and 1.500,
- 2) has high temperature quartz,
- 3) doesn't have biotite.

From these evidence, all of Central part of Kaetsu uplands are decided to M2 terrace and formed on MIS5c.

From this research, I thought that there are precondition for researching about topography of Kaetsu uplands.I think that now, I can research about this area on geomorphology.

Keywords: Kaetsu uplands, marine terrace, MIS5c, tephrochronology

Signatures of ESR signals observed in quartz of Kizu river sediments and in host rocks

Yoshihiko Nosohara^{1*}, Shin Toyoda¹, Masashi Takada², Aiko Shimada³

¹Okayama University of Science, ²Department of History, Sociology and Geography, Faculty of Letters, Nara Women's University, ³JEOL RESONANCE Inc.

Paramagnetic lattice defects in quartz have recently been used as tracers to study the transportation of sediments on the surface of the earth especially for the origin of aeolian dust (e.g. Toyoda and Naruse, 2002), as isotopes have been used. Studies on aeolian dust have used only the E_1' center while impurity centers were shown to be other useful signatures in the studies of river sediments (Shimada, 2008). The present research aims to characterize the sediments of Muro River, a branch river of Kizu River, by analyzing the ESR signals quartz of the sediments. We collected seventeen river sediments and two granites which are possible sources of river sediments. The samples were crushed and sieved to 1000-500, 500-250, and 250-75 micrometers. Quartz grains were extracted from each size fraction by chemical treatment and density separation. The quartz grains were heated at 400 degree celsius for 1 h to erase the inherited ESR signals before gamma ray irradiation. ESR measurements were performed at room temperature and at 81 K to observe E_1' and Ge, and Al and Ti centers, respectively.

The impurity centers were enhanced by gamma ray irradiation. The formation efficiencies of the signals at the origin were obtained, which probably correspond to impurity concentrations in quartz. The E_1' center was observed after heating the sample irradiated to 2.5kGy at 300 degree celsius for 15 min. The formation efficiencies of impurity centers and the intensity of the E_1' center showed constant values in 10 samples of sediments of Muro river with standard deviations of (7 to 12%). It was also found that these values characterize the sediments of each branch river. Details will be reported in the presentation.

Keywords: ESR, quartz, river sediments

Chronology of the terrace surface in the southern part of Kujukuri strand plain, Pacific coast of the Boso peninsula

Kuniyasu Mokudai^{1*}, Futoshi Nanayama², Shinzou Ooi³

¹Pro Natura Foundation Japan, ²Geological Survey of Japan, ³Ibaraki Univ.

The marine terrace around Taito-zaki of the Pacific coast of the Boso Peninsula is considered that formed in Marine Isotope Stage (MIS) 3 by Kuwabara et al. 1999. We surveyed terraces and strata in this area.

Keywords: marine terrace, Ichinomiya River, MIS 3, hill, structural landform

Formation process of the Noshiro coastal dune, northeastern Japan: insight from internal structure and its distribution

Takako Utsugawa^{1*}, SHIRAI, Masaaki¹

¹Graduate School of Urban Environmental Sciences, Tokyo Metropolitan University

The Noshiro coastal dune extends from the Oga Peninsula in a direction NNE through the margin of the northern Hachiro-gata reclaimed land separating from the Japan Sea and is classified as a superpositioning coastal dune. It is about 30 km in length and reaches a height of almost 40-50 m at its peak (the highest peak is 65.2 m in height, Shiraishi 1990). It is composed mainly of detritus originating from the Neogene to the Quaternary strata around the Ou Mountains and delivered by the Yoneshiro River; which has a drainage basin of approximately 4,100 km².

The stratigraphical division; the units composing the Noshiro coastal dune are labeled, in ascending order, "older dunes (Do)", "older humic soils (Ho)", "younger dunes I (Dy-I)", "younger humic soils (Hy)", and "younger dunes II (Dy-II)" and formation processes of the Noshiro coastal dune have mainly been researched by Shiraishi (1990; 1993). However, the following research has not previously been carried out. Based on the characteristic shape of the dune, which is elongated in a longshore (N-S) direction, this study focuses on the formation process of the dune in the longshore direction. We discuss the processes involved in the formation of the dune, based on topographic maps and on the internal structure discussed in our field survey.

A survey using topographic maps and field studies indicated differences between the topographical characteristics and formation processes of the northern and southern parts of the Noshiro coastal dune.

(1) In E-W direction, the northern part (approximately 750 to 1,500 m) of the dune is shown to be wider than the southern part (approximately 500 to 1,000m). (2) The height of the southern parts of the Noshiro coastal dune tends to be higher than that of the northern parts. (3) At the southern site, a wavy humic soil layer was recognized in the lower part of Dy-I. It means that small sand dune bodies had been scattered at Dy-I forming age.

In addition to differences in crustal movements dipping a northeastward, it is evident that there is another factor to be considered to explain the difference between the topographic characteristics. We deduce that a decreasing supply of the beach deposits that supplied the coastal dune sand, caused a change in the formation processes of the northern and southern parts of the Noshiro coastal dune. It is possible, therefore, that swarm of barchans formed a complex internal structure in the southern part of the dune. It would be worth evaluating the sedimentary supply and its affect on the formation process of other coastal dunes situated in a longshore direction.

References

Shiraishi, T. 1990. Holocene geologic development of the Hachiro-gata lagoon, Akita Prefecture, northeast Honshu, Japan. *The Geological Society of Japan* 36: 47-69.

Shiraishi, T. 1993. The Holocene development of the coastal dunes, Akita Prefecture, northeast Honshu, Japan. *The Japanese Geotechnical Society* 41(3): 25-30

Keywords: coastal dune, internal structure, sedimentary supply, barchan, Noshiro region in Akita Prefecture