

The role of tectonics and climate linkage in the Earth System history - An example of the mountain uplift and strengthen

Ryuji Tada^{1*}

¹Graduate School of Science, the University of Tokyo

The linkage between tectonics and climate could be an important process to control earth's surface environment on million years or longer time-scale. Collision of continents and consequent uplift of mountains is an important part of Wilson Cycle during the assembly of super continent and should enhance physical weathering. Enhancement of physical weathering, in turn, increases specific surface area of rocks and minerals in the earth's surface environment and enhance chemical weathering. Enhancement of chemical weathering consumes CO₂ in the atmosphere and also resulted in the increase in nutrients supply to the ocean, the latter further enhance the uptake of CO₂ by the increase in biological productivity in the ocean. Furthermore, formation of supercontinent and/or large plateau will cause strong monsoonal climate that further accelerate chemical weathering through enhanced precipitation. A series of these processes, triggered by continental collision and uplift of mountains, should act as a strong positive feedback to promote global cooling. However, validity of this idea has not been vigorously tested by geological evidence due to the technical difficulty.

Uplift of Himalaya and Tibetan Plateau (HTP) has been believed to have caused enhancement of monsoonal circulation in Asia and pCO₂ decrease during Cenozoic. Because it is the most recent example of the continental collision and regarded as a textbook example of tectonics-climate linkage, it is worth to evaluate the possibility of the positive feedback explained above. Exploring when and how Asian monsoon evolved and whether such evolutionary process was closely related with the uplift of HTP is the first step to test the existence of the positive feedback loop to enhance global cooling. IODP is planning series of expeditions to explore this possibility and Exp. 346 to the Japan Sea and northern East China Sea is the first one to proceed this direction.

Keywords: tectonics, climate, monsoon, chemical weathering, pCO₂, nutrient supply

Quantitative reconstruction of river discharge due to East Asian summer monsoon since the last glacial period in the nor

Yoshimi Kubota^{1*}, Katsunori Kimoto², Ryuji Tada¹, Masao Uchida³, Ken Ikehara⁴

¹University of Tokyo, ²JAMSTEC, ³NIES, ⁴Institute of Geology and Geoinformation, National Institute of Advanced Industrial Science and Techn

In this study, $d^{18}O$ of seawater ($d^{18}O_w$), which is an indirect indicator of sea surface salinity, in the northern ECS during the last 45 ka is reconstructed using Mg/Ca ratio and $d^{18}O$ of planktic foraminiferal shells. According to modern observation, interannual variations in sea surface salinity in summer in the northern part of the ECS is mainly controlled by the discharge from the Changjiang, i.e., rainfall in the drainage area of the Changjiang River. Thus, changes in the sea surface salinity in the northern ECS are interpreted as reflecting variations in the EASM precipitation in South China. It is confirmed that the relationship between salinity in the northern ECS and Changjiang discharge by analyzing the observational salinity data from 1950 to 1994.

The reconstructed freshwater discharge from the Changjiang revealed that there is no long-term decreasing trend in the Changjiang freshwater discharge since the middle Holocene to the present, implying that there is no significant change in EASM precipitation in South China. This result reveals that temporal change in summer precipitation in south China during the Holocene does not follow the summer insolation changes in the northern hemisphere. Instead, millennial-scale variations in the discharge of the Changjiang freshwater are predominant and its variability is larger than decadal variability. The result revealed that variability of the flux of the Changjiang freshwater during the Holocene on centennial to millennial timescale is larger than decadal scale, but much smaller than interannual scale.

On the other hand, based on the $d^{18}O$ balance calculation in this study, it is suggested that surface water in the northern ECS became fresher (1-1.5 PUS lower than present), but estimated average freshwater flux was approximately 25% lower during MIS 3 than during the Holocene. On millennial time scale, lower events of the freshwater discharge coincide with Dansgaard-Oeschger (DO) stadials and Heinrich events in North Atlantic high latitude, while higher discharge events coincide with DO interstadials.

Ice sheet mass balance and the timing of 100,000-year glacial cycles

Ayako Abe-Ouchi^{1*}, Fuyuki SAITO², Kenji Kawamura³, Maureen E. Raymo⁴, Kunio Takahashi², Jun'ichi Okuno², Heinz Blatter⁵

¹AORI, University of Tokyo, ²JAMSTEC, ³NIPR, ⁴Columbia Univ. LDEO, ⁵ETH

The waxing and waning of Northern Hemisphere ice sheets over the past one million years is characterised by an approximately 100,000-year (100-kyr) periodicity and a sawtooth pattern (gradual growth and fast termination){Clark, 2009}{Hays, 1976}. The Milankovitch theory proposes that summer insolation at high northern latitudes drive the glacial cycles{Milankovitch, 1941}, but no significant 100-kyr periodicity exists in insolation intensity{Hays, 1976}. Statistical tests have demonstrated that the glacial cycles are linked to orbital eccentricity, obliquity and precession cycles, presumably through internal feedbacks{Saltzman, 1984}{Tziperman, 2006}{Lisiecki, 2010}{Huybers, 2011}. Furthermore, conceptual models have reproduced the glacial cycles by imposing a threshold of ice volume, ~ excess 100-kyr ice~, for glacial terminations{Raymo, 1997}{Paillard, 1998}{Parrenin, 2003}{Imbrie, 2011}{Huybers, 2011}; however, physical mechanisms have not been identified. Here, using comprehensive climate and ice sheet models, we show that the ~100-kyr periodicity is explained by orbital forcing and internal feedback amongst climate, ice sheet and lithosphere/asthenosphere system. We found that ice sheets exhibit hysteresis responses to summer insolation{Abe-Ouchi, 1993}{Calov, 2005}{Pollard, 2005}, and that the shape and position of the hysteresis loop play a key role to determine the periodicities of glacial cycles. The hysteresis loop of the North American ice sheet is such that, after its inception, the ice sheet mass balance remains mostly positive or neutral through several climatic precession cycles whose amplitude decreases towards eccentricity minimum. The larger it grows and extends towards lower latitudes, the smaller is the required insolation to turn the mass balance to negative. Once the large ice sheet is established, therefore, significantly negative mass balance continues for several thousand years even with a moderate increase in insolation amplitude, allowing time for disintegration. The fast retreat is governed mainly by rapid ablation due to the lowered surface elevation resulting from delayed isostatic rebound. CO₂ plays a supporting, but not controlling, role in the evolution of the ~100-kyr cycle of ice sheet volume and global climate change.

Keywords: climate, climate model, paleoclimate

Paleoclimatic studies using polar ice cores

Kenji Kawamura^{1*}

¹National Institute of Polar Research

Polar ice cores have been extensively used for extracting paleoclimatic and paleoenvironmental information, such as temperature, accumulation, atmospheric aerosols, mineral dust, greenhouse gases, cosmogenic radionuclides, deep-ice biology. They cover the timescales from one year to million years, and the geographical scales from local to global. Here I review the major achievements polar ice core projects, especially those that Japanese communities have played significant roles. They include the Dome Fuji and other ice cores in Antarctica, and NGRIP and NEEM ice cores in Greenland. The future plans of international and Japanese ice core science will also be reviewed.

Keywords: Paleoclimate, Paleoenvironment, Ice core

The Role of the North Pacific in the world ocean circulation

Yusuke Okazaki^{1*}

¹Kyushu University

The North Pacific is known as a terminal region of Ocean Conveyor and abyssal circulation from the south flows into the North Pacific, and upwells to mid-depth and returns south as the Pacific Deep Water (Schmitz, 1996). No deep water forms in the North Pacific in the present ocean because the surface water of the North Pacific is not dense enough to sink into the deep basin (Warren, 1983). Instead, the North Pacific Intermediate Water (NPIW) originated from the Okhotsk Sea lies at depths of 300 to 800 m (Talley, 1993).

The glacial Pacific Ocean had two water masses: well-ventilated and nutrient-depleted glacial North Pacific Intermediate Water (GNPIW) above ~2000 m and less-ventilated and nutrient-enriched deep water below ~2000 m (Keigwin, 1998; Matsumoto et al., 2002). GNPIW is a thicker and more deeply penetrating water mass than the present NPIW. The possible source of GNPIW was possibly in the Bering Sea inferred from microfossil (Ohkushi et al., 2003) and neodymium isotope evidence (Horikawa et al., 2010). A switch of meridional overturning circulation between the North Atlantic and the North Pacific during the last glacial termination was suggested by a collaborative study of sedimentary proxy data and climate modeling (Okazaki et al., 2010).

The North Pacific appears to have played an active role in global ocean circulation, not always passive during the last glacial cycles. The Role of the North Pacific in the world ocean circulation of the past will be discussed.

References:

- Horikawa K, Asahara Y, Yamamoto K and Okazaki Y (2010) Intermediate water formation in the Bering Sea during glacial periods: Evidence from neodymium isotope ratios, *Geology* 38: 435-438.
- Keigwin LD (1998) Glacial-age hydrography of the far northwest Pacific Ocean, *Paleoceanography* 13: 323-339.
- Matsumoto K, Oba T, Lynch-Stieglitz J and Yamamoto H (2002) Interior hydrography and circulation of the glacial Pacific Ocean, *Quaternary Science Reviews* 21: 1693-1704.
- Ohkushi K, Itaki T. and Nemoto N (2003) Last Glacial?Holocene change in intermediate-water ventilation in the Northwestern Pacific, *Quaternary Science Reviews* 22: 1477-1484.
- Okazaki Y et al. (2010) Deepwater Formation in the North Pacific During the Last Glacial Termination, *Science* 329: 200-204.
- Schmitz WJ (1996) Woods Hole Oceanographic Institution Technical Report, WHOI-96-0B, 237 pp.
- Talley LD (1993) Distribution and Formation of North Pacific Intermediate Water, *Journal of Physical Oceanography* 23: 517-537.
- Warren BA (1983) Why is no deep water formed in the North Pacific? *Journal of Marine Research* 41: 327-347.

Significance of the geomorphological approach for the reconstruction of paleo ice-sheets

Hideki Miura^{1*}, OKUNO, Jun'ichi¹, IWASAKI, Shogo², SUGANUMA, Yusuke¹, MAEMOKU, Hideaki³

¹National Institute of Polar Research, ²National University Corporation Tsukuba University of Technology, ³Hiroshima University

In the paleoenvironment research, there are at least two data obtained only by using the geomorphological approach. The one is the history of paleo ice-sheet dimension and subglacial environmental data based on the glacial geomorphological technique, and another is the history of the relative sea-level variations used the coastal geomorphological technique. These data can provide the information about the geographical distribution of past ice sheet, mass balance mechanisms and ice-volume changes using glacial isostatic adjustment model. In this presentation, we introduce the paleo environmental significance of geomorphological approach using the case of East Antarctic ice sheet.

Keywords: ice sheet, glacial geomorphology, subglacial environment, coastal geomorphology, relative sea-level variation, glacial isostatic adjustment

Recent enrichment of nutrient and heavy metal deposition in Japanese mountain lakes due to anthropogenic dust

Narumi Tsugeki^{1*}, Fujio Hyodo², Michinobu Kuwae¹, Wataru Makino³, Yukinori Tani⁴, Jotaro Urabe³

¹Senior Research Fellow Center, Ehime University, ²Research Core for Interdisciplinary Sciences, Okayama University, ³Graduate School of Life Science, Tohoku University, ⁴Institute for Environmental Sciences, University of Shizuoka

Recent studies revealed that nutrient inputs such as nitrogen and phosphorus through atmospheric deposition to aquatic ecosystems have been increasing (Elser et al. 2009; Neff et al. 2008). In addition to nutrients, large emissions of heavy metals such as antimony (Sb) and indium (In) to the atmosphere is expected to be extensive (Filella et al., 2002; Tian et al., 2012; White and Hemond, 2012). East Asia plays an important role in global anthropogenic emissions, but little is known about the effects of nutrient emissions on terrestrial ecosystems and changes in the extent of the metal pollution during its rapid economic growth in recent decades. In this study, we examined fossil pigments and zooplankton remains in dated sediments taken from high mountain lakes at some Japanese National Parks in Hokkaido and Honshu area, to uncover the historical changes of plankton community over the past 100 years. Simultaneously, we measured the geochemical variables such as heavy metals, nitrogen and lead stable isotope to uncover the historical changes of metal deposition, and to identify causal factors including dust source regions. Sedimentary results showed that the fluxes of heavy elements of Sb and In increased at Lake Hachiman-Numa and Hourai-Numa in recent years (Kuwae et al. 2013). Furthermore, the fluxes of phytoplankton abundance in Lake Hourai-Numa (Tsugeki et al. 2012) and Lake Mikurigaike and Niseko-Onuma drastically increased since around 1990 when N stable isotope ratios in sediments decreased, probably due to expanding atmospheric N deposition. In parallel with this, *Daphnia*, a keystone herbivore, increased. During this period, there seems not to be expanding human activities in the watershed around these lakes, suggesting that the increases in nutrients and heavy metals were not resulted from inputs from watershed. Alternatively, Pb stable isotope data in Lake Hachiman-Numa and Hourai-Numa showed that dust deposition with nutrient and metal substances originated from the Asian continent were increasingly transported to study lakes in recent years (Tsugeki et al. 2012; Kuwae et al. 2013). These results imply that long-range transports of anthropogenic dusts have promoted not only Sb and In deposition but also eutrophication in a wide range of Japanese lakes even far from direct human disturbance.

Elser JJ, Andersen T, Baron JS, Bergstrom AK, Jansson M, Kyle M, Nydick KR, Steger L, Hessen DO (2009) Shifts in lake N: P stoichiometry and nutrient limitation driven by atmospheric nitrogen deposition. *Science* 326:835-837.

Filella M, Belzile N, Chen Y-W (2002) Antimony in the environment: a review focused on natural waters: I. Occurrence. *Earth-Science Reviews* 57:125-76.

Kuwae M, Tsugeki NK, Agusa T, Toyoda K, Tani Y, Ueda S, Tanabe S, and Urabe J (2013) Sedimentary records of metal depositions in Japanese alpine lakes for the last 250 years: Recent enrichments of airborne Sb and In in East Asia. *Science of the Total Environment*, 442: 189-197.

Neff JC, Ballantyne AP, Farmer GL, Mahowald NM, Conroy JL, Landry CC, Overpeck JT, Painter TH, Lawrence CR, Reynolds RL (2008) Increasing eolian dust deposition in the western United States linked to human activity. *Nature Geo* 1:189-195.

Tian H, Cheng K, Wang Y, Zhao D, Lu L, Jia W et al. (2012) Temporal and spatial variation characteristics of atmospheric emissions of Cd, Cr, and Pb from coal in China. *Atmospheric Environment* 50:157-63.

Tsugeki NK, Agusa T, Ueda S, Kuwae M, Oda H, Tanabe S, Tani Y, Toyoda K, Wang W, Urabe J (2012) Eutrophication of mountain lakes in Japan due to increasing deposition of anthropogenically-produced dust. *Ecological Research* 27: 1041-1052.

White SJO, Hemond HF (2012) The anthropogeochemical cycle of indium: a review of the natural and anthropogenic cycling of indium in the environment. *Critical Reviews in Environmental Science and Technology* 42:155-186.

Keywords: paleolimnology, Asian continent, anthropogenic dust and nutrient deposition, plankton, heavy metal, stable isotope

High-precision temperature change at the western Japan during the past 3,000 years and its effect on the human activity

Hodaka Kawahata^{1*}, Megumi Matsuoka¹, Ami Togami¹, Naomi Harada², Yusuke Yokoyama¹, Katsunori Kimoto², Yuichiro Tanaka³, Michinobu Kuwae⁴

¹Atmosphere and Ocean Research Institute, The University of Tokyo, ²JAMSTEC, ³AIST, ⁴Ehime University

A continuous record of terrestrial environments is difficult to reconstruct because terrestrial sediments are often eroded and transported away by wind or water. In contrast, marine sediments often provide a continuous record of both marine and terrestrial environments in their sedimentary sequence. Therefore, the continuous reconstruction of paleo-temperature in Holocene was conducted by using coastal marine cores in Hiroshima Bay, which shows very high correlation between the alkenone sea surface temperature (SST) and atmospheric temperature (AT) and enables to estimate a bi-decadal time resolution of record of quantitative temperature in historical period in the western Japan. During the last three millennia, the SSTs (ATs) showed a maximum in 830A.D. (24.3C (25.9C)) and two minima in 780B.C. (22.2C (23.8C)) and 960 A.D. (22.4C (24.0C)) with a mean value of (23.6C (25.2C)), which was comparable to the mean value in mid 20th century natural condition. The low SSTs (ATs) in 800-610B.C. and 990-1190A.D. and relatively low values in 1100-900B.C., 490-320B.C., 550-660A.D. and 1380-1740A.D. corresponded the minima of total solar irradiance (TSI). As TSI change was not sufficient to account for observed temperature amplitude, it is suggested that TSI potentially works as a trigger to drive the other internal forcing in climatic system. Largest volcanic eruptions in 535-536A.D., 1258A.D. and 1452A.D. certainly reduced temperature. Adding instrumental observation records, large shifts in social system such as Hunter-gather to Parry-rice farming and the establishments of Imperial and aristocratic political system, Feudalism and Modern nation, matched large minimal temperatures in cold climate while recovered warm climate afterwards could promote new social systems.

Keywords: Alkenone temperature, Atmospheric temperature, Solar radiation, Historical age, Yayoi People, Japanese

The Pacific Decadal Oscillation and North Pacific regime shifts during the last 2900 years

Masanobu Yamamoto^{1*}, Michinobu Kuwae²

¹Faculty of Environmental Earth Science, Hokkaido University, ²Senior Research Fellow Center, Ehime University

We generated 8-year-resolution records of paleotemperatures using UK37 and the abundance of anchovy and sardine scales in Beppu Bay, Kyushu Island, Japan. Beppu Bay is a small silled basin filled with oxygen-deficient bottom water. Because of oxygen deficient environment, organic matter and fish scales are well preserved in sediments, and bioturbation is limited. Fourteen piston and gravity cores were retrieved at the center of the basin. Correlation of cores was conducted using sand and silt seams (event layers), and the age-depth model was created by wiggle-matching of forty-two AMS radiocarbon dates from bivalve mollusk shells and excess Pb-210 and Cs-137 concentrations. The sedimentation rates were 230-300 cm/ky. UK37 record showed both centennial-scale and multi-decadal variations. Multi-decadal variation, having a significant periodicity at 50-70 years that is typical in the Pacific Decadal Oscillation in the 20th century, is superimposed on centennial-scale variation. The amplitude of the multi-decadal oscillation varied on a multi-centennial scale. The ratio of anchovy to the sum of anchovy and sardine was synchronous with multi-decadal oscillation in SST. Anchovy was more abundant in warmer periods, while sardine was more abundant in cooler periods. This relationship is the same as that was seen in the regime shifts in the 20th century. This is the first evidence showing that North Pacific regime shifts were operated prior to the 20th century.

Keywords: PDO, Regime shift, Holocene, SST, marine core, Beppu Bay

On the possibility of constraining the climate sensitivity: A view from LGM multimodel simulations

Masakazu Yoshimori^{1*}, Ayako Abe-Ouchi¹

¹The University of Tokyo/AORI

Charney climate sensitivity (CCS) is defined as the equilibrium surface temperature change under the doubling of atmospheric CO₂ concentration specifically without vegetation, ice sheet, and carbon cycle feedbacks. CCS is one of the most important metrics in climate projections of coming centuries, and past climate has been used to constrain its uncertainty. The use of paleoclimate as a guide for the future needs to be demonstrated based not only on statistical relation between the past and future climate changes but also on sound physical understanding of mechanisms behind the changes. Much attention has been paid to the last glacial maximum of about 21 thousand years ago, and this presentation overviews previous and current effort on estimating CCS based on LGM climate. The emphasis is placed on the activity with general circulation models and the analysis of the latest PMIP3/CMIP5 multimodels. While perturbed physics ensembles of single models (sensitivity to model parametric uncertainty) suggest a relatively high correlation between LGM and 2xCO₂ global climate feedbacks, multimodel analysis (sensitivity to model structural uncertainty) suggest little correlation between them. This implies that globally averaged LGM climate change does not likely provide a strong constraint on the CCS spread in current models. The radiative feedback analysis indicates that the reason may be cloud feedback induced by the ice sheet forcing unique to the LGM. On the other hand, it was and is proposed that regional change, particularly in the tropics, may be of more use than global mean change. In order to more effectively impose the regional constraint and to increase our confidence, however, uncertainties in proxy data and the forcing estimate need to be reduced and the number of models need to be increased.

Keywords: climate sensitivity, last glacial maximum

Recent progress in researches on biogenic magnetite and applications to paleoceanography

Toshitsugu Yamazaki^{1*}

¹AORI, University of Tokyo

Magnetotactic bacteria, which have chains of magnetite crystals, were discovered in 1975 by Blakemore, and magnetites of bacterial origin preserved in marine sediments (magnetofossil) were first reported in 1986. Since then, biogenic magnetites have been recognized as one of sources of magnetic minerals in sediments. Biogenic magnetites can be identified with TEM from their sizes confined within a single-domain range (several tens of nano-meter) and their characteristic morphologies under biological control. Yet, quantitative estimations such as a proportion of biogenic and terrigenous magnetic minerals were difficult because it is required to extract magnetic minerals for TEM observations, which may distort original magnetic mineral assemblages. However, recent progress of rock magnetic techniques has enabled quantitative estimations for amount and morphology of biogenic magnetites (e.g., Egli, 2010), and it is revealed that biogenic magnetites are dominant magnetic minerals in sediments at least in high latitudes and the equatorial zone (Roberts et al., 2012; Yamazaki and Ikehara, 2012). Quantification of biogenic magnetites have opened new applications to researches on paleoceanography and paleoenvironment, as a kind of fossils. In this presentation, I will introduce examples of such applications.

On the other hand, the discovery for the dominance of biogenic magnetites in sediments has strongly impacted paleomagnetism and its applications, because models of remanent magnetization acquisition processes of sediments did not incorporate contribution of biogenic magnetites. Remanent magnetization of sediments has been explained by a zone magnetization model; remanent magnetization is acquired within a zone with some thickness below the seafloor during compaction after deposition of sediment particles. This implies that there is a lag between a horizon of remanent magnetization fixing and seafloor or the bottom of the bioturbation mixing zone (called lock-in depth). Various models for the amount of the lock-in depth have been presented so far, from few centimeters to more than 40 cm, and have debated for more than thirty years. This is a significant problem when correlating magnetostratigraphy with biostratigraphy and oxygen isotope stratigraphy. Recently, Suganuma et al. (2010) apparently settled the problem by concluding a lock-in depth of ~15 cm from the comparison of geomagnetic paleointensity minimum at the Brunhes-Matuyama polarity transition with abundance of a cosmogenic nuclide ¹⁰Be, which is an independent method for estimating paleointensity devoid of a depth-lag. However, previous arguments on remanent magnetization acquisition mechanism did not seriously consider contribution of biogenic magnetites. It is thought that magnetotactic bacteria live in a sharp chemical gradient from oxic to anoxic in a sediment column and that near the Fe-redox boundary is the most preferable position for magnetotactic bacteria. If this is true, and if biogenic magnetites are the main carrier of remanent magnetization, the magnetization will be fixed near the Fe-redox boundary and the amount of the depth-lag will vary from millimeters to tens of meters depending on sedimentary environments. At present, importance of biogenic magnetites as a carrier of remanent magnetization is not clear even if they are dominant magnetic minerals. This is a significant issue to be solved.

Keywords: biogenic magnetite, magnetotactic bacteria, rock magnetism, paleoceanography, depositional remanent magnetization, Fe-redox boundary

Preliminary study of the Cretaceous tephrochronology in Japan and its application to biostratigraphic study

Reishi Takashima^{1*}, Yuji Orihashi², Toshiro Nagase¹, Sato Kuwabara³, Hiroshi Nishi¹

¹The Center for Academic Resources and Archives, Tohoku University, ²Earthquake Research Institute, The University of Tokyo,

³Faculty of Science, Tohoku University

Tephrochronology is a useful tool for high-resolution stratigraphic correlation and age determination for Quaternary marine and non-marine sediments. However, this method is very difficult to apply to the old sediments such as Paleozoic and Mesozoic because glass in tephra is very sensitive to diagenetic alteration. On the other hand, several heavy minerals in tephra are resistant to diagenetic alteration, and their variations in chemical composition are useful to distinguish individual tephra (Lowe, 2011). Recently, the validity of such mineral analysis as a tool for tephra fingerprinting was supported by Ordovician K-bentonites in North America and Scandinavia (Sell and Samson, 2011).

We analyzed biotite and apatite chemistry of 100-80 Ma tuffs in the Yezo Group exposed in Haboro, Kotanbetsu, Yubari, Hobetsu and Urakawa areas in Hokkaido in order to confirm their efficacy for tephrochronology. Both minerals occur in most tuffs of the Yezo Group. Although some biotites from the lower part of the Yezo Group in Yubari section are mostly altered to chlorite, apatite is always well preserved in all areas and horizons. Binary plot of Mg# vs TiO₂ of biotite and those of MgO vs FeO and F vs Cl of apatite analyses obtained using EPMA are proven to be useful indicators to distinguish individual tuff beds.

Using above method, we identified widely traceable two tuffs in the Yezo Group that are intercalated near the Albian/Cenomanian and the Santonian/Campanian boundaries. These two tuffs from various sections in Hokkaido are plotted in the same field on above mentioned binary plots of biotite and apatite, respectively, and show same U-Pb ages within the margin of error. Correlation of tuffs of the Albian/Cenomanian and the Santonian/Campanian boundaries demonstrate that horizons of first and last occurrences of several age-diagnostic fossils are not always synchronous among areas in Hokkaido. This may attribute to the differences in sedimentary environments and preservation of calcium carbonate among areas. Therefore, tephrochronology using heavy minerals is very useful method to identify "true" biostratigraphic datum, and will improve resolution of biostratigraphy.

References

Sell, B. K. and Samson, S. D., 2011. Apatite phenocryst compositions demonstrate a miscorrelation between the Millbrig and Kinnekulle K-bentonites of North America and Scandinavia. *Geology*, 39, 303-306.

Lowe, D. J., 2011. Tephrochronology and its application: A review. *Quaternary Geochronology*, 6, 107-153.

Keywords: Tephrochronology, Cretaceous, Apatite, Biotite, biostratigraphy

Bio- and chemo-stratigraphy and U-Pb ages of the Cretaceous sequence in Japan

Hiroshi Nishi^{1*}, Reishi Takashima¹, Yuji Orihashi², Toshiro Yamanaka³

¹The Center for the Academic Resources and Archives, Tohoku University, ²Earthquake Research Institute, The University of Tokyo, ³Department of Earth Sciences, Okayama University

Most of GSSPs (Global Boundary Stratotype Section and Points) and their candidate sites of the Cretaceous stages are located in Europe where the stage boundaries are defined by the detailed biostratigraphy (e.g., Gale et al., 1996). Recent studies of integrated stratigraphy of microfossil, megafossil and carbon isotope of those sequences have greatly improved resolution of international stratigraphic correlation. However, few radiometric ages are determined from those sequences because of rare intercalations of volcanoclastic sediments. In order to understand detailed Cretaceous paleo-climatic changes, it is necessary to improve resolution of Cretaceous chronostratigraphic framework based on the radiometric dating.

The Yezo Group, accumulated at approximately 30-40 degree North along the Asian active continental margin in the north-western Pacific Ocean, ranges from Aptian to Paleocene in age and consists mainly of hemipelagic mudstone and turbidite sandstone. Total thickness of this sequence attains 10,000m. This sequence is suitable for establishment of Cretaceous chronostratigraphic framework because it yields abundant felsic tuffs and well preserved age-diagnostic marine macro and micro fossils.

In this study, we established integrated stratigraphies of planktic foraminifera and carbon isotope of wood fragments of the Yezo Group exposed in Tomamae, Yubari and Urakawa sections. Detailed correlation of integrated stratigraphy of planktic foraminifera and carbon isotope between the Yezo Group and European sequences enabled the determination of Cretaceous stage boundaries and oceanic anoxic events (OAEs) in the Yezo Group. The felsic tuffs of the Yezo Group are intercalated at or near stratigraphic datum levels and environmental events, such as Aptian/Albian, Albian/Cenomanian, Cenomanian/Turonian, Turonian/Coniacian, Coniacian/Santonian, Santonian/Campanian boundaries and OAE 1b, OAE1c, OAE1d. The U-Pb ages of these tuff beds are consistent with the inferred age-model of Geologic Time Scale 2012 (GST2012) within the margin of error concerning Aptian/Albian, Albian/Cenomanian, Cenomanian/Turonian, Turonian/Coniacian, Coniacian/Santonian boundaries. On the other hand, Santonian/Campanian boundary, OAE1c and OAE1d show discrepancy about 1 m.y. These discrepancies between this study and GST2012 may attribute to the scarcity of radiometric age of European sections. Therefore, determination of a large quantity of U-Pb age in the Yezo Group will greatly improve the resolution of Cretaceous chronostratigraphy.

References

Gale, A. S., Kennedy, W. J., Burnett, A., Caron, M., Kidd, B. E., 1996. The Late Albian to Early Cenomanian succession at Mont Risou near Rosans (Drome, SE France) : an integrated study (ammonites, inoceramids, planktonic foraminifera, nannofossils, oxygen and carbon isotopes). *Cretaceous Research* 17, 515-606.

Keywords: planktic foraminifer, carbon isotope, Cretaceous, U-Pb age

Biogenic magnetite prevails in oxic pelagic red clay core in the South Pacific Gyre

Takaya Shimono^{1*}, Toshitsugu Yamazaki²

¹Graduate School of Life and Environmental Sciences, University of Tsukuba, ²Atmosphere and Ocean Research Institute, University of Tokyo

Magnetotactic bacteria have been observed in wide variety of environments, including soils, freshwater lakes, and marine sediments, since Blakemore first described in 1975. Magnetotactic bacteria, which most commonly live within the oxic-anoxic transition zone (OATZ) of aquatic environments, produce intracellular crystals of magnetic minerals, specifically magnetite or greigite. It is considered that the magnetite/greigite crystals facilitate the bacteria's search for optimal conditions within the sharp chemical gradients of the OATZ. Petermann and Bleil (1993) reported living magnetotactic bacteria in pelagic and hemipelagic sediments near OATZ in the eastern South Atlantic at water depths to about 3,000 m, but they could not find actively swimming magnetotactic bacteria in sediments of deeper water depths.

The South Pacific Gyre (SPG) is far from continents and the lowest productivity region on Earth. IODP Site U1365 (water depth 5,696 m) cored pelagic red clay of 75.5 m thick above ~100 Ma basement (except for chart layers from ~42 to 61.5 m) in the western edge of the SPG. The core mainly consists of iron rich clay. The color is dark reddish and/or dark brown throughout the core. We conducted a paleomagnetic and environmental rock magnetic study of the pelagic clay core. The magnetostratigraphy revealed the top 5 m sediments cover the last 5 My, and sedimentation rate decreases downward from 1.7 to 0.6 m/m.y. Geochemical measurements of pore water indicate that dissolved oxygen was present throughout the core (>50 microM). Thus oxygen penetrates through the entire sediment column to the sediment/basalt interface, and there is no OATZ.

Magnetic mineral assemblage of this core is dominated by biogenic magnetite despite no OATZ. First-order reversal curve (FORC) diagrams of all specimens have a narrow central ridge along the Hc axis with very small vertical spread. This indicates very weak magnetostatic interaction (Roberts et al., 2000), and is the characteristic of biogenic magnetite (Egli et al., 2010; Roberts et al., 2011). Presence of biogenic magnetite was confirmed by TEM observation. Occurrence of biogenic magnetite was reported also in pelagic red clay of the North Pacific with TEM observations (Yamazaki and Ioka, 1997), and these samples also display the characteristic FORC diagrams. These observations suggest that biogenic magnetites commonly occur in oxic pelagic red clay without OATZ.

Keywords: Biogenic Magnetite, Pelagic Red Clay, Oxic Environment, Environmental Magnetism, South and North Pacific Gyre, IODP Exp.329

Reconstructing the paleoenvironment of the Gulf of Aden and its surroundings lands using biomarkers

Yuta Isaji^{1*}, Hodaka Kawahata¹, Naohiko Ohkouchi², Masafumi MURAYAMA³, Kensaku Tamaki⁴

¹Atmosphere and Oceanic Research Institute, University of Tokyo, ²Japan Agency for Marine-Earth Science and Technology, ³Center for Advanced Marine Core Research, Kochi University, ⁴Graduate School of Engineering, The University of Tokyo

Lands surrounding the Gulf of Aden are one of the most important sites when considering the history of *Homo sapiens*. The oldest known modern human's cranial, estimated to be ca. 195 ka, was found at Omo Kibish, Ethiopia (McDougall et al., 2005). Modern human expansion out of Africa, thought to have taken place at ca. 60 ka (Quintana-Murci et al., 1999), must have been greatly influenced by the environment of the Arabian peninsula. Mesopotamia has seen rises and collapses of many civilizations. The aim of this study is to reconstruct paleoenvironment of these areas at high time resolution. As a first step, I mainly focused on reconstructing paleoceanographic conditions since the ocean has strong interaction with the atmosphere and the land. Here I focused on three periods, 0.7-7.8 ka (Period I), 53-69.7 ka (Period II), and 195-207.4 ka (Period III). Reconstruction of the paleoenvironment was done by extracting lipids from the sediment and analyzing the biomarkers using GC-MS/FID.

The sampling site of the sediment core GOA4 is off the coast of Yemen, in the Gulf of Aden. The climate of the Gulf of Aden is primarily controlled by summer SW monsoon and winter NE monsoon. On a longer time scale, SW monsoon strengthens during the interglacial, and NE monsoon during the glacial (Rostek et al., 1997).

Biomarkers focused in this study are long-chain n-alkanes, alkenones, and highly branched isoprenoids (HBIs). Carbon preference index of the long-chain n-alkanes ranged from 5-8.5, strongly suggesting that they are mainly of terrestrial origin. The long-chain n-alkanes were the only terrestrial biomarker detected in this study. This may imply that the terrestrial environment surrounding the Gulf of Aden had scarce vegetation.

SST reconstructed from core GOA4 was compared with that from core TY93-909/P recovered off eastern Yemen (Rostek et al., 1997). The fact that SST of GOA4 is about 2°C higher during the interglacial indicates that the Gulf of Aden is outside the trajectory of strong SW monsoon. The difference between the SST of GOA4 and TY93-909/P during the interglacial is expected to be larger than the glacial because SW monsoon and following upwelling is more strengthened at the site TY93-909/P. Despite that, fluctuations of SST on glacial-interglacial time scale at both sites show similar trends. Several hypotheses could be made, as follows (Rostek et al., 1997); (i) Global warming (cooling) during the interglacial (glacial) might have cancelled the effect of sea surface cooling (warming) associated with the upwelling strengthening (weakening) of the SW monsoon. (ii) Deepening of the mixed layer due to the enhancement of NE monsoon during the glacial could have counteracted the weakening of the SW monsoon.

The origin of HBIs detected in this study is probably the diatom genus *Rhizosolenia*. Since the size of the genus *Rhizosolenia* is considerably large, the concentration of HBI is used as a proxy for the productivity of diatoms. Sediment trap study taken place in the northwest Arabian Sea indicates that the blooming of diatoms is a month later than that of coccolithophores (Haake et al., 1993). This is because silicate-rich water lies deeper than that of nitrate and phosphate, and injection of silicate-rich water to the surface does not occur until late summer when SW monsoon is more enhanced.

Abundance of diatoms in Period I could be explained by the fact that diatoms prefer nutrient-rich environment. The reason that the coccolithophores were scarce may be because of earlier occurrence of the injection of silicate-rich water to the surface layer due to the enhanced upwelling. This hypothesis does not seem to fit for Period III which scarcity of diatoms can be observed. This may be due to other limiting factors such as Fe availability. Scarce diatoms and abundant coccolithophores in Period II is reasonable since the surface water during the glacial was probably oligotrophic.

Keywords: the Gulf of Aden, human history, biomarker, paleoenvironment, Indian monsoon

Evaluation of the desertification in Tarim Basin based on provenance study of size-separated fluvial sediment since 8Ma

Akinori Karasuda^{1*}, Ryuji Tada¹, Zheng Hongbo², Shin Toyoda³, Hitoshi Hasegawa⁴, Yuko Isozaki¹, Tomohiro Yoshida¹

¹The University of Tokyo, ²Nanjing Normal University, ³Okayama University of Science, ⁴The Nagoya University Museum

Tarim Basin is located to the north of Tibetan Plateau. Tarim Basin is characterized by very dry climate and the Taklimakan desert is developed in the central part of Tarim Basin. In previous studies, the desertification of Tarim basin was considered as having been triggered by the uplift of Tibetan Plateau (e.g. Zheng et al., 2000, DH Sun et al., 2011). However, the timing of desertification has been dated only roughly by eolian sediments, and there is a possibility that Tarim Basin was arid but eolian sediments are not generated.

In this study, we tried to evaluate the timing of aridification based on the dust contribution to the fine fraction in the fluvial sequence which is evaluated by examining provenance of size-separated samples. We conducted the provenance study of quartz, which is common mineral in sediments resistant to weathering.

For the provenance study of quartz, we used Electron Spin Resonance (ESR) signal intensity of quartz and crystallinity index (CI) of quartz. ESR signal intensity of quartz reflects the age of mother rock (Toyoda and Naruse, 2002), whereas CI of quartz reflects physical condition of its formation such as temperature and rate of crystallization (Murata and Norman, 1976). In her study of modern river sediments in the Tarim basin, Isozaki (2009 MS) suggested that quartz in coarse fraction (>63 μ m) of river sediments reflects bedrock geology of the catchment area based on ESR signal intensity and CI of quartz. On the other hand, fine fraction (<16 μ m) of river sediments may reflect geology of the river catchment area and eolian dust.

We applied this method for the fluvial sequence. Firstly, we revealed the size-separated provenance change of fluvial sediments. Secondly, we found the provenance separation of size-separated fluvial sediments, and we evaluated the contribution of eolian dust in the fine fraction of the fluvial sequence.

We conducted field survey at the Yecheng section in the southwestern Tarim Basin. Fluvial to alluvial deposits with occasional intercalations of eolian sediments deposited between 7.6Ma to 1.8 Ma are continuously exposed along the Yecheng section (Zheng et al., 2010; Tada et al., 2010). We selected 27 river sediments (9 sandstones and 18 conglomerate matrix) in 0-16 μ m and 63-500 μ m, and measured ESR signal intensity and CI of quartz.

From the result of analyses, we found the provenance separation of size-separated fluvial sediments (difference in provenance between fine and coarse fractions) at 6.6Ma, 6.0Ma, and after 4.5Ma. Assuming the provenance separation was caused exclusively by eolian contribution, Tarim Basin was arid after at least 6.6Ma.

By comparing with the provenance changes at the Yecheng section in coarse fraction (63-500 μ m)(presented in H-CG33), the ages of uplift activity in the leading edge of Kunlun mountains are approximately the same as the timing of eolian dust contribution increase and desert formation in the Tarim Basin. So, there is possibility that the uplift activity in the leading edge of Kunlun mountains contributed dust production and sand desert formation in the Tarim Basin after 8Ma.

Keywords: Tarim Basin, Desertification, Taklimakan Desert, Provenance study, Eolian dust, Uplift

Estimation of the growth-rate influences on the oxygen isotope and Sr/Ca ratios in the Porites at high latitudes

Shoko Hirabayashi^{1*}, Yusuke Yokoyama¹, Atsushi Suzuki², KAWAKUBO, Yuta¹, Yosuke Miyairi¹, Takashi Okai², NOJIMA, Satoshi³

¹Atmosphere and Ocean Research Institute, The University of Tokyo, ²Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology (AIST), ³Amakusa Marine Biological Laboratory, Kyushu University

The skeletal oxygen isotope in the coral is used as a good proxy to reconstruct sea surface temperature (SST). On the other hand, some studies showed that coral growth rates may affect the oxygen isotope in coral (McConnaughey 1989a; Felis et al., 2003; Suzuki et al., 2005; Hayashi et al., 2013). Here we reanalyzed the same Porites coral in Omata et al. (2006), which was collected in the eastern of Ushibuka in the Amakusa area, Japan, which is located at slightly lower latitude than the northern limit of the hermatypic corals. We measured Sr/Ca ratio and reconstructed SST. Compared to the oxygen isotope ratio measured by Omata et al. (2006), our results showed that Sr/Ca ratio is the robust SST proxy which is independent of its growth rate. We suggest that Sr/Ca ratio is more suitable for reconstruction of SST using the small growth-rate Porites corals, especially in high latitudes.

Keywords: skeletal oxygen isotope in the coral, Sr/Ca ratio, high latitudes, coral growth rates

Provenance and mixing ratio of the sediments discharged from Yangtze River based on ESR signal intensity and Crystallini

Keita Saito^{1*}, Ryuji Tada¹, Tomohisa Irino², Zheng Hongbo³, Chao Luo⁴, Mengying He⁴, Yoshiaki Suzuki¹, Wang Ping⁴

¹EPS, Univ of Tokyo, ²Univ. Hokkaido, ³Nanjing Normal Univ., ⁴Nanjing Univ.

The Yangtze, the largest river in east Asia, is 6300km long and its drainage is 1.94x10⁶km². Since its drainage has over 400 million habitats, the flood of the Yangtze can damage human activities seriously. According to the observation, floods in the middle-lower reaches of the Yangtze are affected by the ENSO. However, floods in the upper reaches are affected by the South Asia Monsoon. So, to investigate floods, not only temporal but also spatial fluctuation is important.

To reconstruct the flood history before the observation records, it is possible to estimate the provenance of flood sediments. For that purpose, it is necessary to establish the proxy to distinguish sediments from each tributaries of Yangtze.

The suspended particle matter, SPM, occupy over 98% of the sediments discharged from Yangtze. Its mean diameter of SPM during normal weather at lower reaches is around 10 μ m. In the flood deposits from the Yangtze estuary, however, the median diameter is 25-35 μ m, which is much larger than that of SPM.

Yang et al., (2007) and Mao et al. (2011) analyzed Sr and Nd isotope in SPM of the Yangtze. CIA is also analyzed by Shao et al. (2012). Their results suggest that it is possible to distinguish sediments from the upper reaches from that in the middle-lower reaches. However, Sr-isotope and CIA can be altered by chemical weathering, and they didn't evaluate qualitative differences depending on particle diameters.

The objective of this study are 1) to distinguish the particles of different tributaries by using ESR(Electron Spin Resonance), CI(Crystallinity Index) and 2) to confirm whether these parameters can estimate the provenance of sediments, which are separated into three fractions.

The result revealed that the ESR values are lower than 2 in the tributaries of the upper reaches, which are gradually rising to 7-10 in the tributaries of the middle reaches. The combination of ESR and CI values can be used to distinguish particles from each tributaries. It is important to select appropriate fractions to analyze, since the ESR values of different fractions are not necessarily the same.

The ESR and CI values at lower reaches are estimated from ESR and CI values of each tributaries and the median sediment budget based on observation. The analyzed values of the lower-reach sediments are slightly different from the estimated values. The sediment budget of each fractions are needed to be improved.

Assuming the sediment flux of tributaries in the upper or middle reaches increases, the magnitudes of changed in ESR and CI values of the sediment from the lower reaches are estimated. These parameters turned out to be more sensitive to the flood of the middle-lower reaches than that of the upper reaches.