

相模湾沿岸域における台風通過に伴う植物プランクトンの一次生産と細菌生産の変動

Typhoon-driven variations of phytoplankton and bacterial production in the coastal waters of Sagami Bay, Japan

土屋 健司^{1*}, 桑原 ピクター 伸一¹, 浜崎 浩二², 多田 雄哉², 今井 章雄³, 市川 忠史⁴, 菊池 知彦⁵, 戸田 龍樹¹

Kenji Tsuchiya^{1*}, KUWAHARA, Shinichi V.¹, HAMASAKI, Koji², TADA, Yuya², IMAI, Akio³, ICHIKAWA, Tadafumi⁴, KIKUCHI, Tomohiko⁵, TODA, Tatsuki¹

¹ 創価大学大学院工学研究科, ² 東京大学大気海洋研究所, ³ 国立環境研究所, ⁴ 水産総合研究センター中央水産研究所, ⁵ 横浜国立大学大学院環境情報研究院

¹ Graduate School of Engineering, Soka University, ² Atmosphere and Ocean Research Institute, The University of Tokyo,

³ National Institute for Environmental Studies, ⁴ National Research Institute of Fisheries Science, ⁵ Graduate School of Environment and Information Science, Yokohama National University

Variations of physical and chemical environments, and responses of phytoplankton and bacterial communities after the passage of typhoon Malou (T1009) in the coastal water of Sagami Bay, Japan, were investigated in order to clarify the effect of a typhoon passage on the temperate coastal ecosystem. Water temperature, salinity, nutrients (dissolved inorganic nitrogen; DIN, phosphate and silicate), dissolved organic carbon (DOC), chlorophyll *a* (Chl *a*), primary production (PP), bacterial abundance (BA) and bacterial production (BP) were measured. Malou approached the study site in Sagami Bay on 8 September 2010 and induced large-scale terrestrial runoff, which led to decrease in salinity to 22.1. Simultaneously with decrease in salinity, concentrations of inorganic nutrients and DOC increased (DIN = 22.4 μM , phosphate = 1.05 μM , silicate = 61.6 μM and DOC = 3.1 mg C L^{-1}), which were more than 3 times higher than the average concentrations of the study site in summer. BP and BP/BA showed relatively fast responses to the passage of Malou reaching maximums of 132 $\text{mg C m}^{-3} \text{d}^{-1}$ and $13.3 \times 10^{-8} \text{ug C cell}^{-1} \text{d}^{-1}$ two days after the passage of Malou, respectively. BA increased gradually reaching a maximum of $2.84 \times 10^9 \text{cell L}^{-1}$ six days after Malou passage. PP/Chl *a* was relatively low just after passage of Malou, and reached a maximum of 184 $\text{mg C} [\text{mg Chl } a]^{-1} \text{d}^{-1}$ three days after typhoon passage. PP showed the highest value of 554 $\text{mg C m}^{-3} \text{d}^{-1}$ five days after Malou passage, and Chl *a* reached a maximum of 7.65 mg m^{-3} six days after typhoon passage. BP and PP showed no correlation and the maximum value of BP/BA was 6.8 times higher than the average value of the study site in summer, which implied that the allochthonous substrate loaded by typhoon passage enhanced bacterial activity just after the passage. BP/PP was highest at 1.5 one day after the passage of Malou, and decreased to 0.1 on the fifth day after typhoon passage suggesting dominant carbon flux changed dynamically from BP to PP in a short time after typhoon passage. Moreover, the integrated PP for 7 days after the passage of Malou accounted for 9% of the annual PP in the upper water of Sagami Bay. Meteorological disturbances such as typhoons, which will likely intensify as global climate change progresses, should be considered in annual production models.

Keywords: typhoon, primary production, bacterial production, temperate coastal region