

中部山岳地域の降雪に含まれる化学成分濃度の空間分布 Spatial distribution of snow chemistry in the Japanese Alps region

倉元 隆之^{1*}, 鈴木 大地², 佐々木 明彦¹, 鈴木 啓助¹

Takayuki KURAMOTO^{1*}, Daichi SUZUKI², Akihiko SASAKI¹, Keisuke Suzuki¹

¹ 信州大学山岳科学総合研究所, ² 信州大学理学部物質循環学科

¹IMS, Shinshu University, ²Dept. Environ. Sci., Shinshu University

The Japanese Alps region is one of the heaviest snowy regions in Japan. In this area winter precipitation is observed mainly two patterns such as winter monsoon pattern and low pressure pattern. Therefore, the chemical characteristics of the snowpack are different by snowfall types. We conducted a snow pit study immediately after snowfall, on the route from Itoigawa, Joetsu and Iida to Matsumoto. We collected only fresh snow samples in winter season. In this study we aimed to clarify spatial distribution of chemical components in fresh snow at the Japanese Alps region. The samples were melted, then pH, electric conductivity and major ions (Na^+ , K^+ , Mg^{2+} , Ca^{2+} , Cl^- , NO_3^- and SO_4^{2-}) were analyzed in clean room. The Na^+ concentration correlates well with Cl^- concentration. These ions are considered to be sea-salt components. On the other hand, SO_4^{2-} concentrations included non-sea-salt components.