

グリーンランド SIGMA-A で観測された積雪粒径と不純物濃度がアルベドに与える効果

Possible effects of snow grain size and snow impurity concentration on the albedo measured at SIGMA-A in Greenland

青木 輝夫^{1*}, 的場 澄人², 山口 悟³, 谷川 朋範⁴, 庭野 匡思¹, 山崎 哲秀⁵, 朽木 勝幸¹, 本山 秀明⁶, 堀 雅裕⁴

Teruo Aoki^{1*}, Sumito Matoba², Satoru Yamaguchi³, Tomonori Tanikawa⁴, Masashi Niwano¹, Tetsuhide Yamasaki⁵, Katsuyuki Kuchiki¹, Hideaki Motoyama⁶, Masahiro Hori⁴

¹ 気象研究所, ² 低温科学研究所, ³ 防災科学技術研究所, ⁴ 宇宙航空研究開発機構, ⁵ 北極探検家, ⁶ 国立極地研究所

¹Meteorological Research Institute, ²Institute of Low Temperature Science, ³National Research Institute for Earth Science and Disaster Prevention, ⁴Japan Aerospace Exploration Agency, ⁵Arctic Explorer, ⁶National Institute of Polar Research

Snow and ice in the Arctic are presently undergoing drastic changes. The snow surface albedo strongly depends on snow grain size and mass concentration of light absorbing impurities. To clarify the contributions of light absorbing snow impurities to recent abrupt melting of snow/ice in Greenland, intensive observations of meteorological and snow parameters have carried out at the site SIGMA-A (78°03'N, 67°38'W, 1,490 m a.s.l.) on northwestern Greenland ice sheet during the intensive observation period (IOP) from June 26 to July 16, 2012. We installed automatic weather station to measure the meteorological elements, radiation budget, snow temperatures, and relative snow height. We have also performed snow pit work and snow samplings for light absorbing snow impurities. During the IOP no precipitation was observed in the first two weeks and a large amount of rainfall with remarkable lowering of snow surface was observed in the middle of July, when a melting event of surface snow/ice over 97% of Greenland ice sheet occurred. Snow grain shapes observed from snow pit work for snow layer of about 80 cm (annual accumulation) changed from melt forms for upper layer and depth hoar beneath that to melt forms for all layers during the IOP. Light absorbing snow impurities found from snow samples were black carbon (BC) and mineral dust, whose concentrations at surface were both increased from 0.9 ppbw to 4.9 ppbw and from 102 ppbw to 1327 ppbw during the IOP, respectively. We calculated the possible albedo reduction by the measured snow impurity concentrations using a physically based snow albedo model. The maximum albedo reduction due even to EC of 4.9 ppbw is less than 0.01 for typical grain size of melt forms, while it could be enhanced by the effect of dust of 1327 ppbw. BC equivalent total impurity concentration is estimated to be about 15 ppbw (albedo reduction -0.015). This result was consistent with the albedos measured during the IOP. The particle size of mineral dust found in surface snow during the latter half period of IOP was larger than 5 micrometers. This indicates a possible transport of mineral dust onto ice sheet surface. Mineral dust is important as nutrient salt for glacial microbial activities, which reduce the albedo in ablation area and could accelerate the melting of ice sheet.

キーワード: グリーンランド, アルベド, 積雪粒径, 光吸収性積雪不純物, 黒色炭素, ダスト

Keywords: Greenland, albedo, snow grain size, light absorbing snow impurities, black carbon, dust