

アラスカのクロトウヒ林における光合成有効放射吸収率の空間依存特性の解析 Spatial scale-dependent characteristics of the fraction of absorbed photosynthetically active radiation

小林 秀樹^{1*}, 鈴木力英¹, 永井信¹, 中井太郎², 金龍元²

Hideki Kobayashi^{1*}, Rikie Suzuki¹, Shin Nagai¹, Taro Nakai², Yongwon Kim²

¹ 海洋研究開発機構, ² アラスカ大学フェアバンクス校国際北極圏研究センター

¹Japan Agency for Marine-Earth Science and Technology, ²International Arctic Research Center, University of Alaska Fairbanks

The fraction of absorbed photosynthetically active radiation (FAPAR) is a ratio of absorbed PAR to incident PAR in plant canopies and is an important vegetation parameter and is widely used for the gross primary production estimation. The absorbed PAR and incident PAR are both sphereradiance (actinic fluxes). The FAPAR definition that ignores the horizontally incident PAR component (FAPAR1d) results in unrealistic FAPAR estimates in heterogeneous forest when looking at high spatial resolution. We investigated the spatial-scale dependence of the relationship between FAPAR1d and the normalized difference vegetation index (NDVI) in highly heterogeneous Alaskan black spruce forest. We collected most of the necessary forest structural datasets used for three-dimensional radiative transfer simulation. At high spatial resolutions (0.1 m), FAPAR1d reaches 6. As the pixels are merged, it converges on the domain-average values. To estimate a domain-average FAPAR from satellite data, 5-meter or coarser resolution is required in sparse forests, depending on the canopy structural conditions and solar geometry.

キーワード: リモートセンシング, 極域, 放射計算

Keywords: remote sensing, polar region, radiative transfer