

流域の栄養塩輸送に及ぼす小規模停滞水域の影響 The effect of small impoundments on nutrient transport in a suburban watershed

清水 裕太^{1*}, 小野寺 真一¹, 大西 晃輝¹, 齋藤 光代², 吉川 昌志¹
Yuta Shimizu^{1*}, Shin-ichi Onodera¹, Koki Onishi¹, Mitsuyo Saito², Masashi Yoshikawa¹

¹ 広島大学大学院総合科学研究科, ² 愛媛大学沿岸環境科学研究センター

¹Graduate School of Integrated and Arts Sciences, Hiroshima University, ²Center for marine environmental studies, Ehime University

There are small impoundments on streams such like weir in the worldwide. In spite of large number of them, the effect of those on nutrients retention is little understood. The objective of this study is to confirm effect of small impoundments on nutrient transport on a catchment scale. Seasonal variation of DN:Cl- ratio which indicates that the ratio increased in summer season and decreased in winter season was confirmed at impoundments. The result of relationship between residence time and difference of concentration of nitrate indicates that greater decrease of nitrate confirmed when it has relatively long residence time. Especially, attenuation of nitrate would start from 2 days of residence time, which is shorter than days that large impoundment start from 8-10 days. Therefore, it is suggested that small impoundments are more effective for nitrate attenuation than large ones. In contrast, the relationship with difference of concentration of DN shows that there is a possibility that the large impoundment could be changed from sink to source. However, small impoundment is easy to maintain itself than large reservoir as dams, by dredging of sediment and removal of colonies of phytoplankton. For conservation of water environment in watersheds which are loaded anthropogenic nitrogen, the small impoundments should be used as places for purification of nitrogen in meaning of a kind of reservoir operation.

キーワード: 小規模貯水池, 滞留時間, 窒素除去, 大和川

Keywords: small impoundment, residence time, nitrogen removal, Yamato River