

GADV peptide/aggregate synthesis using a hydrothermal simulator at elevated temperature
GADV peptide/aggregate synthesis using a hydrothermal simulator at elevated temperature

Kuhan Chandru^{1*}

Kuhan Chandru^{1*}

¹Faculty / Graduate School of Engineering Yokohama National University

¹Faculty / Graduate School of Engineering Yokohama National University

GADV peptide/aggregate synthesis using a hydrothermal simulator at elevated temperature

Kuhan Chandru, Eiichi Imai, Yumiko Obayashi, Takeo Kaneko and Kensei Kobayashi

GADV peptide hypothesis was proposed by Ikehara (2009) as a possible alternative that precedes the RNA world hypothesis (Gilbert, 1986) due to many limitations. The hypothesis suggest a very plausible explanation, but has only been experimentally tested by Oba and co workers (2012), in terms of possible peptide formation using heat-drying cycles and their catalytic activities. We would want to examine the formation of GADV aggregates (or peptides) in an simulated hydrothermal system to represent a more realistic prebiotic environment. We have used Gly, L-Asp, L-Asp and L-Val into our Supercritical Water Flow Reactor (SCWFR) at a temperature range of 100-300 Celsius . Our initial results suggest that no visible aggregates (bigger than 0.2micrometer) were seen after the heating of 2 min in the mentioned temperature range. Initial MALDI-TOF-MS are also suggesting that we only obtained a small peaks about m/z 410 within the spectrum lesser what Oba et al (2012) which was 525, 539, 657. Although many hydrothermal simulation experiments has shown the recovery of amino acids (Islam et al 2002; Kohara et al; Kobayashi et al, 1997) and the formation of oligomers (Imai, 1999; Goto et al , 2005) we only believe, that only small aggregation occur and cannot promote bigger oligomers or polymers due to heat and pressure stress. Hence, based on our initial findings, we are very uncertain about the formation of GADV aggregates or peptides in a hydrothermal system if we use free-form amino acids. It would be of interest to investigate the hypothesis by using bound amino acids or amino acid precursors.

References

Ikehara, K. (2009). Pseudo-Replication of [GADV]-Proteins and Origin of Life. *International Journal of Molecular Sciences*, 10(4), 1525-1537. doi:10.3390/ijms10041525

Gilbert, W (1986) The RNA world. *Nature* 319, 618

Oba, T., Fukushima, J., Maruyama, M., Iwamoto, R., & Ikehara, K. (2005). Catalytic activities of [GADV]-peptides. *Origins of Life and Evolution of the Biosphere : the Journal of the International Society for the Study of the Origin of Life*, 35(5), 447-460.

Kohara, M., Gamo, T., Yanagawa, H., & Kobayashi, K. (2003). Stability of amino acids in simulated hydrothermal vent environments. *Chemistry Letters*, 1-2.

Islam, M. N., Kaneko, T., & Kobayashi, K. (2003). Reaction of Amino Acids in a Supercritical Water-Flow Reactor Simulating Submarine Hydrothermal Systems. *Bulletin of the Chemical Society of Japan*, 76(6), 1171-1178. doi:10.1246/bcsj.76.1171

Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



BAO01-09

会場:301A

時間:5月21日 12:15-12:30

Kobayashi, K., Kohara., Dokiya, M and Yanagawa, H (1997) Formation and stability of amino acids in simulated hydrothermal vent environments. *Viva Origino* 25, 167 - 176

Imai, E.-I., Honda, H., Hatori, K., Brack, A and Matsuno, K (1999) Elongation of Oligopeptides in a Simulated Submarine Hydrothermal System. *Science*, 283(5403), 831-833. doi:10.1126/science.283.5403.831

Goto, T., Futamura, Y., Yamaguchi, Y., & Yamamoto, K. (2005). Condensation reactions of amino acids under hydrothermal conditions with adiabatic expansion cooling. *Journal of Chemical Engineering of Japan*, 38(4), 295-299.

キーワード: GADV, Hydrothermal, prebiotic

Keywords: GADV, Hydrothermal, prebiotic