

南インド、ダルワール超層群における後期太古代のS-MIF化学層序 S-MIF Chemostratigraphy of the Late Archean In the Dharwar Supergroup, South India

三島 郁^{1*}, 山崎里英², Madhusoodhan Satish-Kumar³, 外田 智千⁴, 上野 雄一郎¹
Kaoru Mishima^{1*}, Rie Yamazaki², Satish-Kumar Madhusoodhan³, Tomokazu Hokada⁴, Yuichiro Ueno¹

¹ 東京工業大学, ² 静岡大学, ³ 新潟大学, ⁴ 国立極地研究所

¹Department of Earth and Planetary Sciences, Tokyo Institute of Technology, ²Institute of Geosciences, Shizuoka University,

³Department of Geology, Faculty of Science, Niigata University, ⁴National Institute of Polar Research

Earths tectonic and climatic systems may have fundamentally changed through the late Archean period, which is characterized by major deposition of banded iron formation (BIF) and appearance of stromatolite reef along continental margins. The earliest known glaciation (~2.9Ga) is recorded in the Mozaan Group of South Africa (Young et al., 1998). Also, in the earliest Proterozoic, Snowball Earth event is recorded in the Huronian Supergroup of Ontario, Canada (~2.4Ga; Young et al., 2001). In accord with the climate change, mass-independent fractionation of sulfur isotopes (S-MIF) demonstrated that Earth atmosphere and ocean was oxygenated at around 2.3 Ga from virtually oxygen-free environment (Farquhar et al., 2000). Before the oxidation event, the S-MIF signature changed dramatically: minimum D33S at around 2.9 Ga, subsequent large D33S variation culminated at 2.5 Ga and its sudden drop at the end of Archean. Moreover, D33S-D36S relation shows characteristic ratio of roughly -0.9 in the Archean period. Change of this D36S/D33S relation may reflect the perturbation of atmospheric chemistry. But there is an active debate about the cause of the large D33S variations and D33S-D36S relation through the Archean period.

We studied late Archean volcano-sedimentary sequence of the Dharwar Supergroup, distributed in the Chitradurga Schist Belt, Western Dharwar craton. Our new field mapping and zircon U-Pb dating allows us to reconstruct detailed lithostratigraphy (Hokada et al., 2012). The lower unit (post-3.0 Ga) consists of basal conglomerate, stromatolitic carbonate, silici-clastics with diamictite, chert/BIF and pillowed basalt in ascending order, all of which are older than 2676 Ma magmatic zircon ages from dacite dyke intruded into the topmost pillowed basalt. The upper unit unconformably overlies the pillow lava, and consists of conglomerate/sandstone with ~2600 Ma detrital zircons, komatiite lava, BIF and silici-clastic sequence with mafic volcanics.

Sulfur isotope analysis of extracted sulfide of these sedimentary rocks show a clear MIF and D33S-D36S correlation. The lower group of the Dharwar Supergroup shows D36S/D33S slope of -1.52, middle group shows -1.20, and upper group shows -0.96. This trend is similar to those reported from Pilbara-Kaapvaal equivalents, thus could be a global signature. Moreover, a marked change of D36S/D33S is observed across a diamictite layer (Talya conglomerate) between the lower and the middle group. If this diamictite was glacial in origin, these changes in sulfur isotopes may indicate the link between some transition in atmospheric chemistry and Earth's surface environmental change.

キーワード: 非質量依存同位体分別, 硫黄同位体, ダルワール超層群, 後期太古代, 氷河期

Keywords: mass independent fractionation, sulfur isotope, Dharwar supergroup, late Archean, glaciation