

3.8億年前イスア表成岩: 広域変成作用に基づく太古代沈み込み帯地温勾配、および炭素循環の推定

Regional Metamorphism of the Isua Supracrustal Belt (3.8Ga): Estimate of Archean Geothermal Gradient and Carbon Cycle

新井 達之^{1*}, 大森 聡一², 小宮 剛³, 丸山 茂徳¹

Tatsuyuki Arai^{1*}, Soichi Omori², Tsuyoshi Komiya³, Shigenori Maruyama¹

¹ 東京工業大学大学院地球惑星科学専攻, ² 放送大学文化科学研究科, ³ 東京大学大学院総合文化研究科広域科学専攻

¹Department of Earth and Planetary Sciences, Tokyo Institute of Technology, ²The Faculty of Liberal Arts, The Open University of Japan, ³Department of Earth Science & Astronomy, The University of Tokyo

The 3.7-3.8 Ga Isua Supracrustal Belt (ISB), Southwest Greenland, constitutes the oldest accretionary complex on Earth. Detailed microscopic and microprobe analyses reveal that the west side of ISB comprises metamorphic facies ranging from low to high amphibole facies, which record the Archean geothermal gradient at a subduction zone. Using an isochemical phase diagram (pseudosection), compiled through bulk compositions of ISB, suggests that the geothermal gradient at ISB is an intermediate P/T type in the Archean, whereas high-P/T in Phanerozoic. The shift of the geothermal gradient may reflect the geothermal secular variation of the Earth.

Plate tectonics plays a key role in the carbon global cycling. It has been reported that the less metamorphosed 3.1Ga Archean MORB in Pilbara Craton, West Australia, contain 30 vol% of carbonate minerals in average, formed under the mid-ocean ridge hydrothermal carbonation reaction with the CO₂-rich Archean seawater. On the other hand, the 3.8Ga Archean MORB in the study area, highly metamorphosed under subduction zone, rarely contain carbonate minerals. Comparing the estimated Archean geothermal gradient and stability fields of carbonate minerals of metabasite in the study area, protolith of which is MORB, suggests that most of carbonate minerals in the oceanic crusts cannot be stably dragged into the mantle under the Archean geothermal gradient at the subduction zone even though the oceanic crusts are carbonated up to containing 30vol% of carbonate minerals. Moreover the modal abundance of carbonate minerals in the MORB decreases according to the increasing metamorphic grade ranging from greenschist to middle amphibole facies in the northeast of ISB, which implies that the carbonate minerals must have been formed prior to being subducted at the convergent boundary. Based on these evidences, almost all of carbonate minerals trapped in the oceanic crusts could have returned to the surface at the subduction zone in the Archean even though the Archean oceanic crusts are highly carbonated.

Keywords: Isua Supracrustal Belt, Greenland, Archean, Regional Metamorphism, Geothermal Gradient, Carbon Cycle