

宇宙線生成核種を用いた烏帽子岳周辺における重力性変形地形の発達プロセスの復元

Reconstruction of processes of gravitational slope deformation around Mt. Eboshi using terrestrial cosmogenic nuclide

西井 稜子^{1*}, 松四雄騎², 松崎浩之³

Ryoko Nishii^{1*}, MATSUSHI, Yuki², MATSUZAKI, Hiroyuki³

¹ 筑波大学, ² 京都大学, ³ 東京大学

¹University of Tsukuba, ²Kyoto University, ³The University of Tokyo

A number of linear depressions are widely distributed around Mt. Eboshi, Northern Japanese Alps. This study addressed the reconstruction of formation processes of gravitational slope deformation using terrestrial cosmogenic nuclide. The rock samples for exposure dating were taken from three scarp faces, which are estimated as slip planes related to gravitational slope deformation, corresponding to three linear depressions; two depressions are located at the upper part of the slope and the other is at the middle part of the slope. The concentration of cosmogenic nuclide (¹⁰Be) at the two scarp faces at the upper slope was higher than that at the middle slope. Assuming that initial nuclide concentration is zero and no shielding by seasonal snow cover, we estimated minimum exposure ages of these scarp faces. The minimum exposure ages of upper two scarps were estimated as 2.9 ka and 1.2 ka, while it was calculated to be 0.9 ka for the middle one. These data suggest that slope deformation had advanced downward in Holocene.

キーワード: 宇宙線生成核種, 完新世, 花崗岩類, 北アルプス

Keywords: Terrestrial cosmogenic nuclide, Holocene, granitic rocks, Northern Japanese Alps