

活動的前弧海盆における海底地すべり：東部南海トラフ海域の例 Submarine landslides in an active forearc basin (Eastern Nankai Trough area)

江川 浩輔^{1*}, 伊藤 拓馬¹, 木村 匠¹, 鈴木 清史², 成田 英夫¹
Kosuke Egawa^{1*}, Takuma Ito¹, Sho Kimura¹, Kiyofumi Suzuki², Hideo Narita¹

¹(独)産総研メタンハイドレート研究センター, ²(独)石油天然ガス・金属鉱物資源機構
¹MHRC/AIST, ²JOGMEC

Most of landslide analyses in offshore regions have been spotted in passive continental margins, where glacial-interglacial interaction could activate submarine landslides. These previous studies commonly focus on morphometric feature and distributional pattern of submarine landslide itself, and less on its structural and stratigraphical conditions. Understanding such geological knowledge on landslide regions would be helpful for geohazard assessment and resource development, especially in plate convergent zones. In this meeting, we present seismic and balanced cross sections of mass transport deposits in a Pleistocene forearc basin along the Eastern Nankai Trough to discuss tectono-stratigraphic impact on active-forearc landslides. This study was financially supported by METI and MH21 Research Consortium.

キーワード: 海底地すべり, 前弧海盆, 3次元震探解析, バランス断面法, 東部南海トラフ海域

Keywords: submarine landslide, forearc basin, 3D seismic analysis, balanced cross section, Eastern Nankai Trough area