

## マルチビーム測深地形図の解像度と礁地形の規模

### Resolution of multibeam bathymetric mapping and the dimension of coral reef topography

菅 浩伸<sup>1\*</sup>, 長尾 正之<sup>2</sup>, 中島洋典<sup>3</sup>, 堀 信行<sup>4</sup>, 横山 祐典<sup>5</sup>, 後藤 和久<sup>6</sup>, 大橋倫也<sup>1</sup>, 鈴木 淳<sup>2</sup>, 高田 慎<sup>7</sup>, 中野浩一<sup>7</sup>  
Hironobu Kan<sup>1\*</sup>, Masayuki Nagao<sup>2</sup>, Yosuke Nakashima<sup>3</sup>, Nobuyuki Hori<sup>4</sup>, Yusuke Yokoyama<sup>5</sup>, Kazuhisa Goto<sup>6</sup>, Tomoya Ohashi<sup>1</sup>, Atsushi Suzuki<sup>2</sup>, Shin Takada<sup>7</sup>, Kouichi Nakano<sup>7</sup>

<sup>1</sup> 岡山大, <sup>2</sup> 産総研地質情報, <sup>3</sup> 有明高専, <sup>4</sup> 奈良大, <sup>5</sup> 東大気海洋研, <sup>6</sup> 東北大, <sup>7</sup> (株) 東陽テクニカ

<sup>1</sup> Okayama Univ., <sup>2</sup> Institute of Geology and Geoinformation, AIST, <sup>3</sup> Ariake National College of Technology, <sup>4</sup> Nara Univ.,

<sup>5</sup> AORI, Univ. of Tokyo, <sup>6</sup> IRIDeS, Tohoku Univ., <sup>7</sup> Toyo Corporation

The complex reef topography such as spurs and grooves are difficult to visualize. This study attempts to observe 3D measuring and mapping of outer reef slope using high-resolution multibeam bathymetric sonar. The survey was carried out off the southern coast of Kume Island in the Ryukyu Islands, southwestern Japan. The minimum/maximum depth in the survey area was 0.2/284.7m in the measured area of 1.15 x 1.35 km. The reef topography was visualized with 2 m mesh-size for whole area, 1 m mesh-size for the area shallower than 60 m deep, and 0.2 m mesh-size at ~10 m deep. The bathymetric result was confirmed by SCUBA above the 40 m depth line.

The reef micro-topography (e.g., spurs and grooves) with the dimension ranges from units to tens of meters was visualized when 1 m mesh-size was adopted. The undulation associating with coral colonies which dimension is around tens of centimeters was not visible at the highest resolution map of 0.2 m mesh-size. The reef micro-topography was obscure when the mesh-size larger than 2 m was adopted. This comparison between the map resolution and the topographic dimension is useful for future bathymetric surveys.

Keywords: multibeam bathymetric survey, submarine topography, mapping, scale, coral reef, Kume Island