Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



会場:301B

時間:5月19日15:45-16:00

ALOS/AVNIR-2とPRISMを用いた水災害に関する脆弱性評価データの生成 Generation of Vulnerability Assessment data about Water-related Disaster using ALOS/AVNIR-2 and PRISM

中郡 俊文^{1*}, 田林雄², 桑原 祐史³, 齋藤修⁴, 安原一哉⁵ Toshifumi Nakagori^{1*}, TABAYASHI, Yu², Yuji Kuwahara³, SAITO, Osamu⁴, YASUHARA, Kazuya⁵

¹ 茨城大学理工学研究科,² 筑波大学陸域環境研究センター,³ 茨城大学広域水圏環境科学教育研究センター,⁴ 茨城大学防 災セキュリティー技術教育研究センター,⁵ 茨城大学地球変動適応科学研究機関

¹Graduate School of Science and Engineering, Ibaraki University, ²Terrestrial Environment Research Center, Tsukuba University, ³Center for Water Environment Studies, Ibaraki University, ⁴Center for Intellectual Disaster Prevention Studies, Ibaraki University, ⁵Institute for Global Change Adaptation Science, Ibaraki University

According to the Fourth Assessment Report of the IPCC, due to the climate change, it is assumed that the flood damages such as typhoon will be increased at lowland area. In order to decrease storm and flood damages, it is important to unify the measures over land use, environmental plan, and traffic environment. In this research, we focused on the Hai Hau coast that is a downstream of Red-River, Vietnam. Around the Hai Hau coast, vast lowland areas are distributed. Therefore, in this research, we mapped the vulnerability assessment to the river flood for Hai Hau coast and its vicinity. Then, we mapped a salt farm, a shrimp farm and cultivated land using ALOS/AVNIR-2 and ALOS/PRISM in order to grasp the vulnerability assessment of the cultivated land received from overtopping waves of the coast. The result of this study, we generated the geographic information for water-related disaster, and considered the relation with present condition of land-use.

キーワード: 気候変動, 衛星画像, 氾濫シミュレーション, リモートセンシング Keywords: climate change, satellite image, flood simulation, remote sensing