

海洋堆積物深部のアミノ酸の生物地球化学的動態：化合物レベル窒素同位体組成とD/L比からの制約

Biogeochemical dynamics of amino acids in deep-subsurface marine sediments

山口 保彦^{1*}, 力石 嘉人², 高野 淑識², 小川 奈々子², 菅 寿美², 横山 祐典³, 大河内 直彦²

Yasuhiko T. Yamaguchi^{1*}, Yoshito Chikaraishi², Yoshinori Takano², Nanako O. Ogawa², Hisami Suga², Yusuke Yokoyama³, Naohiko Ohkouchi²

¹ 東京大学大学院理学系研究科, ² 海洋研究開発機構 海洋・極限環境生物圏領域, ³ 東京大学大気海洋研究所

¹Department of Earth and Planetary Science, The University of Tokyo, ²Institute of Biogeosciences, JAMSTEC, ³Atmosphere and Ocean Research Institute, The University of Tokyo

Amino acids in sediment pore waters are key compounds in metabolic activities of sedimentary microbes and in remineralization of carbon and nitrogen. However little is known about their biogeochemical dynamics (e.g., sources and transformation processes) in deep-subsurface sediments.

As a new approach to constrain the sources of dissolved amino acids in sediment pore waters, this study reports and compares compound-specific d15N and enantiomer ratio (%D) of total hydrolysable amino acids (THAA) in solid phase and dissolved hydrolysable amino acids (DHAA) in pore waters from the same sediment samples. Samples were collected from deep-subsurface sediments (down to 172.9 m below seafloor) at the Sagami Trough (NW Pacific) during D/V Chikyu cruise CK09-03 (Expedition 905: December 2009).

In the sediments deeper than 9 mbsf, average %D values of DHAA were 25.9% in alanine, 24.8% in aspartic acid, 11.3% in serine, and 16.3% in glutamic acid, and average %D changes from THAA were +15.3% in alanine, -0.4% in aspartic acid, -8.1% in serine, and 4.6% in glutamic acid. Compound-specific d15N analysis showed that d15N values of alanine are higher in the DHAA than the THAA and that d15N values of glycine and glutamic acid are similar between the two fractions (d15N-DHAA - d15N-THAA = +5.8 permil, +1.9 permil, -0.3 permil, respectively). These results suggest that the DHAA fractions have different d15N and %D signatures from the THAA fractions, and that hydrolysis of the THAA could not be the sole source of the DHAA. Alternatively, the d15N and %D signatures of DHAA are consistent with the idea that in situ release of proteinaceous materials from sedimentary microbial biomass (such as peptidoglycan of Gram-positive bacteria) is an important source of DHAA. This suggests that recycle of dissolved amino acids by microbes would be an important process during amino-acid degradation and microbial metabolism in the deep-subsurface sediments.

キーワード: 海底下生命圏, 有機物, 窒素同位体, アミノ酸, バクテリア

Keywords: Deep biosphere, Organic matter, Nitrogen isotope, Amino acids, Bacteria