

## サハリン沖 LV59 堆積物コア間隙水の化学分析 Chemical analysis of LV59 sediment pore waters retrieved off Sakhalin Island, Russia

南尚嗣<sup>1\*</sup>, 八久保 晶弘<sup>1</sup>, 坂上 寛敏<sup>1</sup>, 久保 圭佑<sup>1</sup>, 小竹 毅<sup>1</sup>, 山下 聡<sup>1</sup>, 高橋 信夫<sup>1</sup>, 庄子 仁<sup>1</sup>, Young K. Jin<sup>2</sup>, Boris Baranov<sup>3</sup>, Anatoly Obzhairov<sup>4</sup>

Hirotsugu Minami<sup>1\*</sup>, Akihiro Hachikubo<sup>1</sup>, Hirotsushi Sakagami<sup>1</sup>, Keisuke Kubo<sup>1</sup>, Tsuyoshi Kotake<sup>1</sup>, Satoshi Yamashita<sup>1</sup>, Nobuo Takahashi<sup>1</sup>, Hitoshi Shoji<sup>1</sup>, Young K. Jin<sup>2</sup>, Boris Baranov<sup>3</sup>, Anatoly Obzhairov<sup>4</sup>

<sup>1</sup> 北見工業大学, <sup>2</sup> 韓国海洋研究開発研究所, <sup>3</sup> ロシア科学アカデミー P.P. シルシヨフ海洋学研究所, <sup>4</sup> ロシア科学アカデミー極東支部 V.I. イリチェフ太平洋海洋学研究所

<sup>1</sup>Kitami Institute of Technology, <sup>2</sup>Korea Polar Research Institute, <sup>3</sup>P.P. Shirshov Institute of Oceanology RAS, <sup>4</sup>V.I. Il'ichev Pacific Oceanological Institute FEB RAS

The Sakhalin Slope Gas Hydrate Project (SSGH) is an international collaborative effort by scientists from Japan, Korea, and Russia to investigate natural gas hydrates (GHs) that have accumulated on the continental slope off Sakhalin Island, Okhotsk Sea. In 2012, the field operation of the SSGH-12 project was conducted as the LV59 cruise of the R/V Academic M.A. Lavrentyev. GH-bearing and -free sediment cores were retrieved using steel hydro- and gravity corers.

The pore water was sampled on board using squeezers designed and constructed at the Kitami Institute of Technology (KIT). A 10-cm depth interval of the sediment core was drawn into the squeezer, and pore water was then directly collected into a polyethylene syringe connected to the discharge tube of the squeezer. The other end of the syringe was connected to a membrane filter cartridge containing a 0.2-um filter. Seawater samples were obtained from the top of the corer. All water sample was filtrated through a 0.2-um filter and then taken into a polypropylene bottle.

The concentrations of anions in the pore waters were measured at KIT. Sulfate concentrations decreased linearly with depth to the sulfate methane interface (SMI). The linearity of these profiles suggested that sulfate depletion is largely driven by an upward flux of methane, rather than by the flux of organic matter from above, and the anaerobic oxidation of methane (AOM) at the SMI. The linear sulfate profiles are formed in a steady state based on sulfate and methane co-consumption at the SMI and the balance of sulfate and methane fluxes. The SMI of the GH-bearing LV59-27HC core was 0.5 m below the seafloor (m bsf). Since the depth of the SMI depends on the intensity of the upward methane flux, we can conclude that intensive methane flux was observed at the GH-bearing core.

キーワード: ガスハイドレート, 堆積物間隙水, 化学分析, ロシアサハリン島  
Keywords: gas hydrate, pore water, chemical analysis, Sakhalin Island, Russia