

極域下部熱圏中性風の研究への期待 Study of the lower thermospheric wind dynamics

野澤 悟徳^{1*}
Satonori Nozawa^{1*}

¹ 名古屋大学太陽地球環境研究所
¹STEL, Nagoya University

The lower thermospheric wind dynamics has been paid attention for several decades to understand the Magnetosphere-Ionosphere-Thermosphere coupling. In particular, it has been an issue how the lower thermosphere will response to the solar wind energy input. IS radar measurements of the polar lower thermosphere begun about 40 years ago by a pioneer work of Brekke et al. [JGR, 78, 8235, 1973], and significant number of studies have been published since then. However, our understanding of the lower thermosphere is still limited. One of reasons is that the lower thermosphere is significantly influenced by atmospheric waves propagating from below. Thus, the day-to-day variability is very prominent. Owing to high running cost, long run of IS radar had not been conducted a decade ago. In 2007-2008, EISCAT Svalbard radar was operated almost continuously for 1 year. However, only about 20 % of the data sets can be used for deriving the ion velocity vector. If we have wind velocity datasets on daily basis like meteor and MF radars usually made for the mesospheric wind measurements, our understanding of the lower thermosphere wind dynamics will be much more progressed. EISCAT 3D can make it possible. I will overview works of the lower thermospheric wind and propose what kind of run we desire.

キーワード: EISCAT 3D, 下部熱圏, 北極域
Keywords: EISCAT 3D, Lower thermosphere, polar region