

EISCAT_3Dを用いたイオン上昇流/流出の研究 Study on ion upflow and outflow with EISCAT_3D

小川 泰信^{1*}
Yasunobu Ogawa^{1*}

¹ 国立極地研究所
¹ National Institute of Polar Research

An important phenomenon of magnetosphere-ionosphere coupling is the formation of upwelling ions in the topside polar ionosphere. These upflows can be a significant loss of atmospheric gasses into interplanetary space and a significant source of magnetospheric plasma, which may also affect the dynamics of the magnetosphere. Key processes for upward ion flows in the topside ionosphere are suggested to be frictional heating, ambipolar diffusion driven by a heated electron gas, and transverse ion acceleration produced by plasma waves. It is critical to determine the relative importance of the different mechanisms in operation and to understand the 3D distribution and composition of the upflowing ions and neutrals. Moreover, there are several transitions of upflowing ions, for examples, from chemical to diffusion dominance at 500-800 km altitude, from subsonic to supersonic flow at 1000-2000 km altitude, and from collisional to collisionless region at 1500-2500km altitude. EISCAT_3D is one of the most suitable measurements to investigate such transitions because of its wider height coverage (up to about 2000 km) along the field line. EISCAT_3D will have more transmitter power density and higher sensitivity than those of the current Tromso UHF radar, and will give information of accurate thermal ion velocity, upward flux, and ion composition (O⁺, H⁺, and hopefully NO⁺). In this paper, potential investigations of ion upflow and outflow using the EISCAT_3D are shown, and desirable specifications of the EISCAT_3D are discussed.

キーワード: 極域電離圏, EISCAT_3D, イオン流出
Keywords: Polar ionosphere, EISCAT_3D, ion outflow