

## 樹木年輪中 $^{14}\text{C}$ 濃度にみられる突発イベント Rapid events in the carbon-14 content of tree-ring

三宅 芙沙<sup>1\*</sup>, 増田 公明<sup>1</sup>, 中村 俊夫<sup>2</sup>  
Fusa Miyake<sup>1\*</sup>, Kimiaki Masuda<sup>1</sup>, Toshio Nakamura<sup>2</sup>

<sup>1</sup> 名古屋大学太陽地球環境研究所, <sup>2</sup> 名古屋大学年代測定総合研究センター

<sup>1</sup>Solar-Terrestrial Environment Laboratory, Nagoya University, <sup>2</sup>Center for Chronological Research, Nagoya University

Measurement of cosmogenic nuclides, which are radioisotopes produced by cosmic rays, can provide us important information to search past extraterrestrial high-energy events such as supernova, solar proton event (SPE), and so on. Until now, the contents of  $^{14}\text{C}$  in tree-rings and  $^{10}\text{Be}$  in ice cores have been used for this purpose. However, no clear evidence has been found by  $^{14}\text{C}$  and  $^{10}\text{Be}$ .

We show the results of  $^{14}\text{C}$  content measurement in Japanese cedar annual rings from AD 600 to 1020 with 1- to 2-year resolution, and report two findings of rapid increases of  $^{14}\text{C}$  content from AD 774 to 775 and AD 992 to 993. These are clear increases against its measurement errors. The shapes of the two series are very similar, i.e., a rapid increase within one year followed by a decay due to the carbon cycle. The scale of the AD 993 event is 0.6 times as large as the AD 775 event.

The  $^{10}\text{Be}$  flux in the Antarctic ice core also shows peaks corresponding to these two  $^{14}\text{C}$  events. The proportions of flux increase ( $^{14}\text{C}/^{10}\text{Be}$ ) of the two events are consistent with each other. Therefore, it is highly possible that these events have the same origin.

Although the cause of this event can be explained by a large solar proton event (SPE) or a short gamma-ray burst, we conclude that solar activity is a plausible cause because the occurrence rate of  $^{14}\text{C}$  increase events is inconsistent with a observed rate of an short gamma-ray bursts.