

## 磁場なし太陽風での磁気圏電離圏対流 The magnetosphere-ionosphere convection in the solar wind without magnetic fields

藤田 茂<sup>1\*</sup>, 片岡 龍峰<sup>2</sup>, 細川 敬祐<sup>3</sup>, 田中 高史<sup>4</sup>  
Shigeru Fujita<sup>1\*</sup>, Ryuho Kataoka<sup>2</sup>, Keisuke Hosokawa<sup>3</sup>, Takashi Tanaka<sup>4</sup>

<sup>1</sup> 気象大学校, <sup>2</sup> 東京工業大学, <sup>3</sup> 電気通信大学, <sup>4</sup> 九州大学

<sup>1</sup>Meteorological College, <sup>2</sup>Tokyo Institute of Technology, <sup>3</sup>University of Electro-communication, <sup>4</sup>Kyushu University

We investigate the magnetosphere-ionosphere convection when merging between the interplanetary magnetic field (IMF) and the magnetospheric one is not effective. To perform the simulation in this condition, we assign the IMF with extremely small intensity. The simulation reveals that the magnetosphere has long magnetotail unlike the closed magnetosphere proposed by Axford and Hines [1961]; this means traditional viscous interaction may not play an important role. It is also obtained that the ionospheric convection exhibits a 2-cell pattern like the magnetosphere-ionosphere convection system for the southward IMF condition. It is revealed that the magnetosphere-ionosphere convection is associated with the R1 field-aligned current generated in the cusp-mantle region where the convection plasma flow traverses the cusp region with enhanced plasma pressure. The traversing plasma flow in the cusp region is fed by the magnetosheath flow which enters into the cusp region along the magnetic field lines. Thus, the magnetosphere-ionosphere stationary convection is performed without the merging.

キーワード: 極端宇宙天気, 太陽風-磁気圏相互作用, 磁気圏電離圏対流, MHD シミュレーション, 複合系, 粘性相互作用  
Keywords: Extreme space weather, solar wind-magnetosphere interaction, magnetosphere-ionosphere convection, MHD simulation, compound system, viscous interaction