Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.





PEM05-35

会場:303

## 太陽高エネルギー粒子被ばく線量予測システム WASAVIES の開発:システムの概要 Development of WASAVIES (Warning System of AVIation Exposure to SEP): System Overview

佐藤 達彦<sup>1\*</sup>, 保田 浩志<sup>2</sup>, 片岡 龍峰<sup>3</sup>, 八代 誠司<sup>4</sup>, 桑原 孝夫<sup>5</sup>, 塩田 大幸<sup>6</sup>, 久保 勇樹<sup>7</sup> Tatsuhiko Sato<sup>1\*</sup>, Yasuda Hiroshi<sup>2</sup>, Ryuho Kataoka<sup>3</sup>, Seiji Yashiro<sup>4</sup>, Takao Kuwabara<sup>5</sup>, Daiko Shiota<sup>6</sup>, Yuki Kubo<sup>7</sup>

<sup>1</sup>日本原子力研究開発機構,<sup>2</sup>放射線医学総合研究所,<sup>3</sup>東京工業大学,<sup>4</sup>アメリカカトリック大学,<sup>5</sup>デラウェア大学,<sup>6</sup>理 化学研究所,<sup>7</sup>情報通信研究機構

<sup>1</sup>JAEA, <sup>2</sup>NIRS, <sup>3</sup>TITech, <sup>4</sup>Catholic University of America, <sup>5</sup>University of Delaware, <sup>6</sup>RIKEN, <sup>7</sup>NICT

When solar energetic particles (SEP) are incident to the atmosphere, they can induce air showers by generating varieties of secondary particles. Such secondary particles can reach deep into the atmosphere, and enhance the level of radiation doses, which can be a hazard of aircrews. In order to precisely estimate the radiation doses during large SEP events, we are developing a warning system of aviation exposure to SEP, WASAVIES. In the system, the time profile of anisotropic SEP spectrum incident to the Earth are estimated from the latest knowledge of space weather information [1,2], such as basic parameters of flare and coronal mass ejections (CME). The enhancement of the radiation doses at flight altitudes during the event is then calculated from the incident SEP fluxes in combination with a database developed based on air shower simulation performed by the PHITS code [3]. The WASAVIES has been tested and verified by making a comparison between the measured and calculated count rates of several neutron monitors during past GLE (ground level enhancement) events. The final goal of our project is to predict the enhancement of radiation doses due to SEP exposure within 6 hours from the GLE onset.

[1] Kataoka et al. JpGU2013

[2] Kubo et al. JpGU2013

[3] http://phits.jaea.go.jp/

キーワード: 太陽高エネルギー粒子, 宇宙線被ばく, 空気シャワーシミュレーション, WASAVIES, PHITS Keywords: SEP, Cosmic-ray exposure, Airshower simulation, WASAVIES, PHITS