

サブオーロラ帯における地上 VLF 波動観測 - VLF-CHAIN キャンペーン Ground-based VLF wave observations at subauroral latitudes - VLF-CHAIN Campaign

塩川 和夫^{1*}, 横山 侑¹, 家田 章正¹, 三好 由純¹, 野村 麗子¹, 李 星恩¹, 尾崎光紀², 石坂和大², 砂川尚貴², 八木谷 聡², 片岡 龍峰³, 土屋 史紀⁴, イアン・スコフィールド⁵, マーチン・コナーズ⁵
Kazuo Shiokawa^{1*}, Yu Yokoyama¹, Akimasa Ieda¹, Yoshizumi Miyoshi¹, Reiko Nomura¹, Sungeun Lee¹, Mitsunori Ozaki², Kazumasa Ishizaka², Naoki Sunagawa², Satoshi Yagitani², Ryuho Kataoka³, Fuminori Tsuchiya⁴, Ian Schofield⁵, Martin Connors⁵

¹ 名古屋大学太陽地球環境研究所, ² 金沢大学, ³ 東京工業大学, ⁴ 東北大学, ⁵ アサバスカ大学

¹Solar-Terrestrial Environment Laboratory, Nagoya University, ²Kanazawa University, ³Tokyo Institute of Technology, ⁴Tohoku University, ⁵Athabasca University

We report observations of very low frequency (VLF) waves during the VLF Campaign observation with High-resolution Aurora Imaging Network (VLF-Chain) of February 17-25, 2012, at subauroral latitudes at Athabasca (54.72N, 246.69E, MLAT=61.3). Continuous measurements of VLF waves with a sampling rate of 100 kHz have been made since then to monitor daily variations of chorus waves and their detailed structures. We found quasi-periodic (Q-P) emissions for which their repetition period rapidly changes within one hour without corresponding magnetic pulsations and for which their intensity suddenly increased associated with a storm sudden commencement without changing their frequency. Patchy burst in the upper-band frequency ranges are often observed during magnetically disturbed times. Falling tone chorus whose rate of frequency change varies on a timescale less than a minute was observed. Clear systematic correlation of these various chorus waves with cosmic noise absorption was not seen throughout the campaign period. These observations indicate existence of several new types of VLF wave phenomena at subauroral latitudes.

キーワード: VLF 波動, サブオーロラ帯, 地上観測

Keywords: VLF wave, subauroral latitudes, ground-based observation