Japan Geoscience Union Meeting 2013

(May 19-24 2013 at Makuhari, Chiba, Japan)

©2013. Japan Geoscience Union. All Rights Reserved.



会場:101B



時間:5月20日16:45-17:00

Observation of the O2 (0-0) atmospheric band nightglow by the IMAP/VISI: a case study Observation of the O2 (0-0) atmospheric band nightglow by the IMAP/VISI: a case study

Septi Perwitasari^{1*}, Takeshi Sakanoi¹, Atsushi Yamazaki², Yuichi Otsuka³, Yuta Hozumi⁴, Yusuke Akiya⁴, Akinori Saito⁴ Septi Perwitasari^{1*}, Takeshi Sakanoi¹, Atsushi Yamazaki², Yuichi Otsuka³, Yuta Hozumi⁴, Yusuke Akiya⁴, Akinori Saito⁴

¹PPARC Tohoku University, ²JAXA/ISAS, ³STEL Nagoya University, ⁴Geophysics Dept. Kyoto University ¹PPARC Tohoku University, ²JAXA/ISAS, ³STEL Nagoya University, ⁴Geophysics Dept. Kyoto University

The Visible and near-Infrared Spectral Imager (VISI) of the IMAP mission has been launched onto the International Space Station (ISS) since July 21st 2012. IMAP/VISI begins its nominal operation from the middle of October and will make a continuous observation for the next 3 years. IMAP/VISI is now operated in the nightside hemisphere with a range of +/- 51 deg. GLAT, measuring three different airglow emissions of OI at 630 nm, the OH Meinel band at 730 nm and the O2 (0-0) atmospheric band at 762 nm at an altitude of ~400 km with the typical spatial resolution of 16 ? 50 km. Since the start of nominal operation in the middle of October 2012, IMAP/VISI has been observing airglow emission in the MLT region and found many interesting features. One of the rare events observed by IMAP/VISI is a concentric pattern of gravity wave on the O2 (0-0) airglow emission. This pattern was observed on October 18, 2012 over northeastern part of Japan around 1200 UT. The similar pattern was also observed by all-sky camera at Rikubetsu in the 557.5 nm and OH airglow emissions. From the MTSAT satellite, we found a strong convective activity over Honshu Island around the same hour, which could be the source of this rare pattern. We will report the case study of this event and a possibility to do the statistic study on this concentric pattern if we could find other events from IMAP/VISI observation.

 $\neq - \nabla - F$: IMAP/VISI, O2 (0-0) atmospheric band, nightglow, concentric gravity wave structu Keywords: IMAP/VISI, O2 (0-0) atmospheric band, nightglow, concentric gravity wave structu