

## 航空機からのハイスピードカメラ観測により明らかとなった スプライトストリーマの時間・空間進展機構 Spatial and Temporal Evolution of Sprite Streamers Derived from High-Speed Camera Data in Aircraft Observation Campaign

小林 縫<sup>1\*</sup>, 佐藤 光輝<sup>1</sup>, 高橋 幸弘<sup>1</sup>, 工藤 剛史<sup>1</sup>, 三宮 佑介<sup>1</sup>, 井上智広<sup>2</sup>, H. C. Stenbaek-Nielsen<sup>3</sup>, Matthew G McHarg<sup>4</sup>, Ryan K Haaland<sup>5</sup>, Takeshi Kammae<sup>3</sup>, Yoav Yair<sup>6</sup>, Walter A Lyons<sup>7</sup>, Steven A Cummer<sup>8</sup>, NHK Cosmic Shore Project<sup>2</sup>  
Nui Kobayashi<sup>1\*</sup>, Mitsuteru Sato<sup>1</sup>, Yukihiko Takahashi<sup>1</sup>, Takeshi Kudo<sup>1</sup>, Yusuke Sanmiya<sup>1</sup>, Tomohiro Inoue<sup>2</sup>, H. C. Stenbaek-Nielsen<sup>3</sup>, Matthew G McHarg<sup>4</sup>, Ryan K Haaland<sup>5</sup>, Takeshi Kammae<sup>3</sup>, Yoav Yair<sup>6</sup>, Walter A Lyons<sup>7</sup>, Steven A Cummer<sup>8</sup>, NHK Cosmic Shore Project<sup>2</sup>

<sup>1</sup>北海道大学理学院宇宙理学専攻, <sup>2</sup>NHK制作局 宇宙の渚プロジェクト, <sup>3</sup>Univ. of Alaska, Fairbanks, AK, United States, <sup>4</sup>US Airforce Academy, Colorado Springs, CO, United States, <sup>5</sup>Fort Lewis College, Durango, CO, United States, <sup>6</sup>Open University, Ra'anana, Israel, <sup>7</sup>FMA Research, Ft Collins, CO, United States, <sup>8</sup>Duke University, Durham, NC, United States

<sup>1</sup>Department of Cosmospaces, Hokkaido Univ., Sapporo, Japan, <sup>2</sup>Japan Broadcasting Corp. NHK Cosmic Shore Project, Tokyo, Japan, <sup>3</sup>Univ. of Alaska, Fairbanks, AK, United States, <sup>4</sup>US Airforce Academy, Colorado Springs, CO, United States, <sup>5</sup>Fort Lewis College, Durango, CO, United States, <sup>6</sup>Open University, Ra'anana, Israel, <sup>7</sup>FMA Research, Ft Collins, CO, United States, <sup>8</sup>Duke University, Durham, NC, United States

The occurrence conditions of sprites streamers still remain to be an unsolved problem after the discovery of sprites. Though the detailed three-dimensional spatial structures and the temporal evolution of sprite streamers are the key parameters to clarify the occurrence conditions, these spatiotemporal characteristics are not clearly identified. In order to specify the detailed spatial and temporal evolution of sprite streamers, we have conducted the optical observation campaign using high-speed cameras from two jet aircrafts in summer US. In this campaign, we succeeded to capture sprite images for 28 events by the high-speed cameras with a sampling rate over 8,000 fps at each aircraft simultaneously. Using these image data, we have performed triangulation analysis to estimate the horizontal spatial distribution and vertical extent of sprite streamers. We have analyzed two types of columniform sprites; one is the columniform sprite with a preceding dense inhomogeneous halo, and the other is the columniform sprite with a preceding dim halo or without a halo. In the later case (dim halo plus columns), the following results are identified. (1) The longer the distance between sprite columns and the parent CG becomes, the higher the bottom altitude of columns becomes. (2) The longer the distance between sprite columns and the parent CG becomes, the slower the speed of downward streamer tips becomes. These results are first clear observational evident showing the horizontal spatial gradient of the quasi-electrostatic field produced by the parent CG discharge. At the presentation, we will show the electrical characteristics of the parent CG discharges derived from CMC waveforms and will discuss the possible mechanisms determining such spatial dependences.