

径方向に開いた太陽風中の非線形アルヴェン乱流によるイオン加熱 Ion heating by the nonlinear evolution of low-frequency Alfvénic turbulence in the radially expanding solar wind

成行 泰裕^{1*}, 梅田隆行², 鈴木建³, 羽田亨⁴

Yasuhiro Nariyuki^{1*}, Takayuki Umeda², Takeru K. Suzuki³, Tohru Hada⁴

¹ 富山大・人間発達, ² 名大・太陽地球環境研究所, ³ 名大・理学研究科, ⁴ 九大・総理工

¹Univ. Toyama, ²STEL, Nagoya Univ., ³Department of Physics, Nagoya Univ., ⁴ESST, Kyushu Univ.

It is well known that the magnetohydrodynamic (MHD) turbulence with the high cross helicity (Alfvénic turbulence) is ubiquitously observed in the solar wind plasmas. The Alfvénic turbulence generated in the vicinity of the photosphere may play an important role in heating of coronal plasmas, and a number of authors have investigated the "wave modeling" of the solar wind plasmas. On the other hand, while the heating of ions is dominated by the collisionless damping of the low-frequency waves, few models including the low-frequency Alfvénic turbulence have gone beyond the MHD description. In this study, we present the nonlinear evolution of low-frequency Alfvénic turbulence in the radially expanding solar wind by using a kinetic-fluid model (Vlasov-MHD model). The heating of ions by low-frequency Alfvénic turbulence in the absence of the ion cyclotron resonance is demonstrated. In order to discuss the self-consistency of the ion kinetics, we carry out the test-particle simulation, in which the numerical data of the global MHD simulation is used. The other non-MHD effects are also discussed as the phenomenological parameters.

キーワード: 太陽風, アルヴェン乱流, イオン加熱

Keywords: solar wind, Alfvénic turbulence, ion heating