

衝突過程でのトロクトライトにおけるオリビンと斜長石の組織変化 The change of texture of olivine and plagioclase by the shock process in troctolite

武藤 大樹^{1*}, 鹿山 雅裕², 関根 利守³
Daiki Mutou^{1*}, Masahiro KAYAMA², Toshimori Sekine³

¹ 広島大学大学院地球惑星システム学専攻, ² 広島大学大学院理学研究科地球惑星システム学専攻, ³ 広島大学大学院理学研究科

¹Department of Earth and Planetary Systems Science, Graduate School, Hiroshima University, ²Department of Earth and Planetary Systems Science, Graduate School of Science, Hiroshima University, ³Graduate School of Science, Hiroshima University

The collision phenomena are important for understanding evolution of planets, their surfaces. Although there are a lot of experimental studies on plagioclase and olivine, two constituent minerals of an ordinary meteorite, we need to know the shock wave effects on these minerals more detailed. We conducted shock experiments on a troctolite using a single stage propellant gun at NIMS in Tsukuba. At peak pressures of 10 GPa, 20 GPa, 30 GPa, 45.5 GPa and 58.3 GPa. The recovered samples are investigated by SEM, EPMA, XRD and the cathode luminescence method (CL method). In samples recovered at 10 GPa and 20 GPa, the intense cracks were observed. Above 30 GPa, however, the polished surface of plagioclase became, indicating that it transformed into glass. At 45.5 GPa and 58.3 GPa, the textures indicated partial melting. From XRD analyses, we found that the diffraction peaks of plagioclase decreased greatly at 30 GPa and most plagioclases changed amorphous. The CL emission intensity decrease. There are three peaks centered at about 330 nm, 400 nm and 550 nm for the initial plagioclase. In the 30 GPa sample, two peaks centered at ~ 330 nm and 380 nm. The EPMA analyses indicates no compositional change but a slight deviation from the total of 100 %.

On the other hand, the irregular cracks were observed in olivines recovered samples from 10 GPa, 20 GPa, 30 GPa and 45.5 GPa. At 58.4GPa, however, cracks were not observed, suggesting that olivine became soften.

During analysis by EPMA, amphiboles were found in all samples except the sample at 20 GPa, and calcite left at 58.4 GPa. On the other hand, clay minerals disappeared above 45.5 GPa. We keep detailed study further.

キーワード: トロクトライト, 衝撃圧力, 衝撃実験, オリビン, 斜長石
Keywords: troctolite, shock pressure, shock experiment, olivine, plagioclase