

ハイパースペクトルリモートセンシングによる月スピネル全球搜索 Global Survey of Lunar Spinel-Rich Exposures by Satellite Hyperspectral Remote Sensing

山本 聡^{1*}, 中村 良介², 松永 恒雄¹, 小川 佳子³, 石原 吉明², 諸田 智克⁴, 平田 成³, 大竹 真紀子⁵, 廣井 孝弘⁶, 横田 康弘¹, 春山 純⁵

Satoru Yamamoto^{1*}, Ryosuke Nakamura², Tsuneo Matsunaga¹, Yoshiko Ogawa³, Yoshiaki Ishihara², Tomokatsu Morota⁴, Naru Hirata³, Makiko Ohtake⁵, Takahiro Hiroi⁶, Yasuhiro Yokota¹, Junichi Haruyama⁵

¹ 国環研, ² 産総研, ³ 会津大, ⁴ 名大, ⁵ JAXA/ISAS, ⁶ ブラウン大

¹NIES, ²AIST, ³Univ. of Aizu, ⁴Nagoya Univ., ⁵JAXA/ISAS, ⁶Brown Univ.

The studies using the spectral data obtained by Spectral Profiler (SP) and Multiband Imager (MI) onboard the Japanese lunar explorer SELENE/Kaguya revealed the global distributions of the purest anorthosite (PAN), olivine-rich materials, and orthopyroxene-rich materials over the entire Moon. These results were based on the diagnostic bands of these lunar major minerals in spectral data with wavelength less than 1.7 micron. Recently, a prominent Mg-spinel-rich material (hereafter, Mg-spinel) on the lunar surface has been identified by Moon Mineralogy Mapper onboard Chandrayaan-1. Since the Mg-spinel is characterized by a strong absorption band around 2 micron, the spectral data with wavelength longer than 1.7 micron are needed to find the Mg-spinel by remote-sensing spectral data. We have recently updated the radiometric calibration for SP NIR 2 data with wavelength longer than 1.7 micron. Based on the entire data set of SP, including the SP NIR 2 data, we conducted the global survey to find the Mg-spinel on the Moon. Here, we report the global distribution of the Mg-spinel sites based on this survey.

キーワード: リモートセンシング, ハイパースペクトル

Keywords: remote-sensing, hyperspectral